



Effect of Sulphur and Bio-inoculants on Growth and Yield of Soybean (*Glycine max* L)

Jason Masih ^{a++*}, A. A. Kawade ^{a#} and V. V. Panchal ^{a#}

^a Shri Vaishnav Institute of Agriculture, SVVV, Indore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/acri/2024/v24i121025>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128662>

Original Research Article

Received: 20/10/2024

Accepted: 24/12/2024

Published: 29/12/2024

ABSTRACT

An experiment entitled, "Effect of sulphur and bio-inoculants on growth and yield of soybean (*Glycine Max* L.)" was conducted during *kharif*, 2023 at new experimental cum demonstration field, SVIAG, SVVV, Indore. The field experiment was carried out in randomized block design with nine treatments. The treatments were T₁- RDF (20:40:20), T₂- RDF + 20 kg S ha⁻¹, T₃- RDF + 20 kg S ha⁻¹ + *Rhizobium*, T₄- RDF + 20 kg S ha⁻¹+ PSB, T₅- RDF + 20 kg S ha⁻¹+ *Thiobacillus*, T₆- RDF + 20 kg S ha⁻¹ + *Rhizobium* + PSB, T₇- RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus*, T₈- RDF + 20 kg S ha⁻¹ + *Thiobacillus* + PSB and T₉- RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus* + PSB. The growth and yield attributes such as plant height, no. of leaves plant⁻¹ at 60 DAS, no. of branches plant⁻¹, leaf area plant⁻¹ at 60 DAS, dry matter plant⁻¹, number of pods plant⁻¹, pod yield plant⁻¹, seed yield plant⁻¹, seed yield (ha⁻¹), straw yield (ha⁻¹) and biological yield (ha⁻¹) were recorded and found

⁺⁺M. Sc Agronomy;

[#] Assist. Prof of Agronomy;

*Corresponding author: Email: jasonmasih03@gmail.com;

Cite as: Masih, Jason, A. A. Kawade, and V. V. Panchal. 2024. "Effect of Sulphur and Bio-Inoculants on Growth and Yield of Soybean (*Glycine Max* L)". Archives of Current Research International 24 (12):342-46. <https://doi.org/10.9734/acri/2024/v24i121025>.

to be highest with the application of RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus* + PSB as compared to rest of the treatments. Application of RDF along with sulphur (20 kg ha⁻¹) and seed treated with all bio-inoculants (*Rhizobium*, PSB and *Thiobacillus*) or at least one of the available bio-inoculant helps to increase soybean growth and yield of soybean.

Keywords: Sulphur; bio-inoculants; rhizobium.

1. INTRODUCTION

The soybean (*Glycine max*) is a species of legume native to East Asia and belongs to the Fabaceae or bean family. Beans are classed as pulses whereas, soybeans are classed as oilseeds. Due to its high protein content, soybean is known as "poor man's meat". It contains 18-20 per cent oil and 40 per cent protein. The oil and protein content together account for about 60 per cent of dry soybeans by weight. The remainder consists of 35 per cent carbohydrate and about 5 per cent ash. Soybean protein is rich in valuable amino acid, lysine (5 %) which is deficient in most cereals. It also contains 60 per cent polyunsaturated fatty acids (52.8 % linolenic acid + 7.2 % linoleic acid). Soybean has high caloric value releasing 432 calories from 100 gm edible protein as compared to 350 calories from cereals of the same quantity. The oil contains about 0.5-1.0 per cent lecithin which is essential for the building up of human nerve tissues.

"As per world soybean production, United States was the leading global producer of soybeans with a production volume of 120.52 million metric tons in 2018-19. As of 2019, Brazil overtook the United States as the leading soybean-producing country with a production volume of some 162 million metric tons in 2022-23 followed by United States of America, Argentina, China and India accounting for 40 per cent of world production. In India, the area, production and productivity of soybean during 2022-23 was 12.07 m ha, 13.98 Mt and 1158 kg ha⁻¹ respectively as against 12.18 m ha area, 13.12 Mt production and 1077 kg ha⁻¹ productivity during 2021-22. Among the states, Madhya Pradesh stood first with 50.18 lakh ha followed by Maharashtra, Rajasthan, Karnataka, Gujarat and Telangana". (Anonymous 2023).

"Recent years have seen the use of chemical fertilizers in agriculture, making the nation more self-sufficient in food production, but at the expense of the ecosystem and the well-being of all living things, creating negative impacts on soil fertility. To satisfy our agricultural requirements, beneficial microorganisms are better alternatives

to conventional farming methods. Bioinoculants do not show any detrimental impact on the soil, plant and animal life as they are eco-friendly and highly efficient. Nitrogen-fixing bacteria, such as *Rhizobium*, form intimate alliances with soybean roots, establishing nodules where they convert atmospheric nitrogen into a plant-usable form. This biological nitrogen fixation not only reduces the need for synthetic fertilizers but also enhances soil fertility, enriching the nutrient bank through enhancing nutrient availability and uptake for present and subsequent crops and promoting sustainable soil management practices, the key drivers of soybean growth and yield. Phosphorus-solubilizing bacteria (PSB) unlock phosphorus in the soil, making it accessible to soybean roots and fuelling critical metabolic processes. *Thiobacillus*, a genus of sulphur-oxidizing bacteria, plays a significant role in the sulphur cycle in soil environments. These bacteria facilitate the conversion of sulphur and sulphide compounds into forms that plants can readily absorb i.e. sulphate. The activity of *Thiobacillus* can enhance soil fertility by increasing a steady supply of sulphate thus, leading to better crop growth and higher yields" (Shahwar et al., 2023).

The global demand for soybean continues to rise due to its nutritional benefits and use in various industrial applications. This increasing demand necessitates advancements in cultivation practices to enhance yield and ensure sustainability. The soybean crop's adaptability to different environmental conditions and its ability to enhance soil fertility through nitrogen fixation makes it an integral component of sustainable agricultural systems. The idea of combining sulphur fertilization with the bio-inoculant application can create a synergistic effect that promotes sustainable soybean production. Sulphur will enhance the efficiency of biological nitrogen fixation by supporting the enzymatic processes involved, while bio-inoculants improve the availability and uptake of nutrients, including sulphur and phosphorus. Thus, the research work to study the effect of sulphur and bio-inoculants on soybean production was selected and the results obtained are discussed herewith.

2. MATERIALS AND METHODS

The present experiment entitled “Effect of sulphur and bio-inoculants on soybean production (*Glycine max* L.)” was carried out at New experimental cum demonstration field, Shri Vaishnav Institute of Agriculture, Shri Vaishnav Vidyapeeth Vishwavidyalaya, Indore during *rabi* 2023-24. The soil at the experimental site was clayey in texture, low in available nitrogen (206.18 kg ha⁻¹), medium in phosphorus (13.98 kg ha⁻¹), and high in available potassium (448.25 kg ha⁻¹). The field experiment was carried out in randomized block design with nine treatments and three replications. The treatments were T₁- RDF (20:40:20), T₂- RDF + 20 kg S ha⁻¹, T₃- RDF + 20 kg S ha⁻¹ + *Rhizobium*, T₄- RDF + 20 kg S ha⁻¹ + PSB, T₅- RDF + 20 kg S ha⁻¹ + *Thiobacillus*, T₆- RDF + 20 kg S ha⁻¹ + *Rhizobium* + PSB, T₇- RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus*, T₈- RDF + 20 kg S ha⁻¹ + *Thiobacillus* + PSB and T₉- RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus* + PSB. The field was divided into 27 plots with gross plot size of 3.00 m x 4.50 m each.

Recommended dose of fertilizers (20:40:20 NPK kg ha⁻¹) and sulphur was applied as basal dose and seed treatment with different bio-fertilizers (*Rhizobium*, *Thiobacillus* and PSB) was given as per treatments. The nutrients were applied through Urea, Single super phosphate (SSP), Muriate of potash (MOP) and Bensulph (Elemental Sulphur) as basal dose before sowing.

“To record biometric observations, five plants were chosen at random from each net plot. Seedlings emerged six days after sowing, and the total number of plants was counted 15 days later. The final plant stand was recorded at the time of harvesting. Height of plant were measured from ground level to the base of last fully opened leaf and average height was work out. The total number of trifoliolate leaves were counted and recorded. Number of branches plant⁻¹ grown on stem were counted and recorded. Leaf area were calculated by using formula. A single plant was randomly sampled from each plot for dry matter production. Numbers of pods were counted from 5 randomly selected plants from net plots and then the average was worked out. Total numbers of pods from five selected sample plants were harvested and per plant seed yield was recorded. Thousand seeds were counted from each net plot seed yield and its weight was recorded. The

plants from each net plot were harvested and seeds were separated by threshing. After sun drying seed yield obtained in each net plot were weighed in kg and presented as kg ha⁻¹. Straw yield was obtained by deducting the seed yield from the weight of total dry produce (biological yield) of respective net plot in kg and given as kg ha⁻¹. The figures of biological yield were calculated by summing of seed yield and straw yield of net plots. Finally, it was converted on hectare basis. The harvest index was calculated by using formula seed yield (kg ha⁻¹) ÷ biological yield (kg ha⁻¹) X 100” (A.A. Kawade et al.; 2016).

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

The growth parameters such as plant height increased at every step of crop development till maturity. The number of branches increased up to 60 days and remained constant thereafter. The number of nodules increased positively up to flowering and decreased thereafter due to shrinkage of nodules. The number of leaves and leaf area increased up to 60 days and decreased thereafter towards maturity due to leaf senescence. Total dry matter increased at every growth stage from 30 DAS up to harvest. The development of pods started after flowering and increases continuously up to harvest. Highest values of plant height (42.53 cm) at harvest, no. of leaves plant⁻¹ (32.87) at 60 DAS, no. of branches plant⁻¹ (12.13) at harvest, leaf area plant⁻¹ (13.40) at 60 DAS and dry matter plant⁻¹ (23.53 g) at harvest was recorded in treatment with application of RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus* + PSB (T₉) inoculants throughout the entire crop growth period. The lowest values of all these parameters were recorded by the application of RDF alone (T₁). The increase in growth attributes may be due to better uptake and translocation of plant nutrients to growing plants. The findings were supported by Jaga and Sharma (2015), Farhad et al., (2010), Akter (2013), Kawde et al., (2016), Aziz et al., (2016) and Sharma (2011).

3.2 Yield Attributes and Yield

Yield attributing characters viz., maximum no. of pods plant⁻¹ (31.90), pod yield plant⁻¹ (22.67 g), seed yield plant⁻¹ (11.33 g) at harvest was recorded in treatment with the application of RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus* + PSB (T₉). Whereas the minimum values were recorded from the treatment with the application of RDF alone (T₁).

Table 1. Effect of sulphur and bio inoculants on growth attributes of soybean

Treatments	Plant height (at harvest)	No. of leaves plant ⁻¹ (60 DAS)	No. of branches plant ⁻¹ (at harvest)	Leaf area plant ⁻¹ (60 DAS)	Dry matter plant ⁻¹ (at harvest)
T ₁	33.70	24.33	7.12	7.33	12.22
T ₂	33.88	26.33	8.53	7.48	17.68
T ₃	35.87	27.41	9.82	9.59	19.50
T ₄	34.07	27.03	9.56	8.79	18.20
T ₅	36.56	27.47	9.92	10.37	20.02
T ₆	39.37	29.47	10.99	12.19	21.95
T ₇	40.53	30.20	11.82	12.28	22.31
T ₈	38.85	28.93	10.63	11.60	21.26
T ₉	42.53	32.87	12.13	13.40	23.53
S.Em. (±)	1.78	0.85	0.59	0.51	1.02
CD at 5%	5.35	2.56	1.77	1.53	3.06
General mean	37.26	28.23	10.06	10.34	19.63

Table 2. Effect of sulphur and bio inoculants on yield attributes of soybean

Treatments	No. of pods plant ⁻¹	Pod yield (g) plant ⁻¹	Grain yield (g) plant ⁻¹	Seed yield q ha ⁻¹	Straw yield q ha ⁻¹	Biological yield q ha ⁻¹
T ₁	21.03	13.17	7.67	8.11	13.99	22.09
T ₂	21.57	15.23	7.83	10.27	17.22	27.49
T ₃	24.20	18.20	8.40	12.11	21.58	33.69
T ₄	22.73	17.00	7.97	11.76	20.28	32.04
T ₅	26.87	19.43	9.17	13.21	24.45	37.66
T ₆	30.00	21.07	10.10	15.08	27.10	42.18
T ₇	30.47	21.43	10.67	15.92	27.87	43.80
T ₈	29.13	20.07	9.87	14.21	25.94	40.15
T ₉	31.90	22.67	11.33	16.41	28.52	44.93
S.Em. (±)	0.93	0.88	0.56	1.06	0.92	1.48
CD at 5%	2.79	2.65	1.68	3.18	2.76	4.44
General mean	26.43	18.70	9.22	13.01	22.99	36.00

The seed, straw and biological yield of maize was significantly influenced by different treatments. Application of RDF + 20 kg S ha⁻¹ + *Rhizobium* + Thiobacillus + PSB (T9) recorded maximum seed yield (16.41 q ha⁻¹), straw yield (28.52 q ha⁻¹) and biological yield (44.93 q ha⁻¹) which was found to be at par with RDF + 20 kg S ha⁻¹ + *Rhizobium* + PSB (T6), RDF + 20 kg S ha⁻¹ + *Rhizobium* + Thiobacillus (T7) and RDF + 20 kg S ha⁻¹ + Thiobacillus + PSB (T8). The increase in all the growth parameters with application of RDF + 20 kg S ha⁻¹ + *Rhizobium* +

Thiobacillus + PSB (T9) must have resulted in increase in final yield of soybean. The increase in all the growth and yield attributes in T9 treatment must have resulted in higher seed, straw and biological yield. These findings are supported by Bonde and Gawande (2017) and Patel et al., (2013).

4. CONCLUSION

It may conclude that; the highest values of growth attributes, yield attributes and yield can

be obtained with the application of RDF + 20 kg S ha⁻¹ + *Rhizobium* + *Thiobacillus* + PSB (T9) whose values are at par with the treatment containing any one of the bio-inoculant.

Application of RDF along with sulphur (20 kg ha⁻¹) and seed treated with all bio-inoculants (*Rhizobium*, PSB and *Thiobacillus*) or at least one of the available bio-inoculant helps to increase soybean growth and yield of soybean.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Akter, F. M. D., Islam, N., Shamsuddoha, A. T. M., Bhuiyan, M. S. I., & Shilpi, S. (2013). Effect of phosphorus and sulphur on growth and yield of soybean (*Glycine max* L.). *International Journal of Bio-resource and Stress Management*, 4(4), 555-560.
- Anonymous. (2023). The soybean outlook.
- Aziz, A. L. A., Ahiabor, B. D. K., Opoku, A., & Abaidoo, R. C. (2016). Contributions of *Rhizobium* inoculants and phosphorus fertilizer to biological nitrogen fixation, growth, and grain yield of three soybean varieties on a fluvic luvisol. *American Journal of Experimental Agriculture*, 10(2), 1-11.
- Bonde, A. S., & Gawande, S. N. (2017). Effect of integrated nutrient management on growth, yield, and nutrient uptake by soybean (*Glycine max*). *Annals of Plant and Soil Research*, 19(2), 154-158.
- Farhad, I. S. M., Islam, M. N., Hoque, S., & Bhuiyan, M. S. I. (2010). Role of potassium and sulphur on the growth, yield, and oil content of soybean (*Glycine max* L.). *Academic Journal of Plant Sciences*, 3(2), 99-103.
- Jaga, P. K., & Sharma, S. (2015). Effect of bio-fertilizer and fertilizers on productivity of soybean. *Annals of Plant and Soil Research*, 17(2), 171-174.
- Kawade, A. A., Panchal, V. V., & Kalegore, N. K. (2016). Influence of sulphur and bio-inoculants on growth and yield of black gram (*Vigna mungo* L.). *National Academy of Agricultural Science*, 34(7), 2113-2117.
- Kawde, A. A., Panchal, V. V., & Kalegore, N. K. (2016). Influence of sulphur and bio-inoculants on yield and economics of black gram (*Vigna mungo* L.). *National Academy of Agricultural Science*, 34(7), 2119-2122.
- Patel, H. R., Patel, H. F., Maheriya, V. D., & Dodia, I. N. (2013). Response of kharif green gram to sulphur and phosphorus fertilization with and without biofertilizer application. *International Journal of Life Sciences*, 8(1), 149-152.
- Shahwar, D., Mushtaq, Z., Mushtaq, H., Abdulaziz, A., Alqarawi, P. Y., Thobayet, S., & Alshahrani, F. S. (2023). Role of microbial inoculants as biofertilizers for improving crop productivity: A review. *Heliyon*, 9, 2405-8440.
- Sharma, A. (2011). Economic viability of soybean production with sulphur and phosphorus nutrients on yield and yield attributes of soybean. *Journal of Progressive Agriculture*, 2(2), 13-15.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/128662>