



Status of Rice Diseases in Northern Himalayas: A Case Study of Himachal Pradesh

**Shikha Sharma ^{a*}, DK Banyal ^a, Sonali Parwan ^{a*},
Shiwali Dhiman ^b, Sachin Upmanyu ^c, Neelam Bhardwaj ^d
and Anchal Thakur ^a**

^a Department of Plant Pathology, College of Agriculture, CSKHPKV, Palampur, India.

^b KVK Dhaulakuan, Sirmour, CSKHPKV, Palampur, India.

^c Department of Plant Sciences, Central University of Himachal Pradesh, Dharamshala, India.

^d Department of Genetics and Plant Breeding, College of Agriculture, CSKHPKV, Palampur, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author SS conducted surveys, lab studies and collected seed samples as well as recorded data. Authors DKB, SD and SU collected seed samples and analysed the data. Author SP conducted lab studies and drafted the manuscript. Authors NB and AT managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/acri/2025/v25i11031>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128744>

Short Research Article

Received: 28/10/2024

Accepted: 30/12/2024

Published: 02/01/2025

ABSTRACT

Rice production around the world is threatened by various diseases leading to biotic stress. Surveys during 2022 & 2023 were conducted in eight districts of Himachal Pradesh to assess the status of rice diseases. Seed health status of farmers saved seed was also monitored. Surveys of the crop

*Corresponding author: Email: shikhasharmapp@hillagric.ac.in, parwansonali1610@gmail.com;

revealed the occurrence and incidence of varying degrees of diseases viz., bacterial leaf blight, neck blast, sheath rot, false smut, brown spot, narrow brown spot, rice stunt and leaf scald. Association of mycoflora i.e *Ustilaginoidea virens*, *Pyricularia oryzae* and *Helminthosporium oryzae* in farmer saved seed was also observed. The results of this study can be implemented to curtail the yield losses associated with seed borne diseases and can further improve sustainable rice production in Himachal Pradesh.

Keywords: Seed health; pathogens; diseases; rice; incidence.

1. INTRODUCTION

Rice (*Oryza sativa* L.) is considered as one of the major staple cereal food source for more than half of the world's population and forms a significant component of the energy resource of the human race. According to Udemezue (2018), rice is the world's second most important cereal crop following corn and also one of the food crops mostly grown in tropical and sub-tropical regions of the world. About 522.65 million metric tons (Mmt) of milled rice are produced worldwide each year with 137.83 Mmt being produced in India (USDA, 2024). Rice is an economically and scientifically important crop which is mostly produced and consumed in Asia and African countries (Dede et al. 2019; Islam et al. 2000). With more than 40% of global rice exports, India plays a pivotal role in ensuring food security across the globe. The assessment of seed health standard in rice is very important for farmers and food security as it is a first line approach in managing seed-borne diseases. Furthermore, the quality of planted seeds has a critical influence on the ability of crops to become established and to realize their full yield potential and value (McGee, 1995). Seed-borne diseases may not only introduce new pathogens to affect the quantity or quality of the crop yield but also contaminate the soil permanently (Anselme, 1981). Most crop diseases that are important economically are seed-borne and seed transmitted, including blast disease of rice, bakanae, loose smut, flag smut, karnal bunt, and ear cockle of wheat (Akter & Hossain, 2015). Worldwide, nearly 56 fungi pathogens are reported to infest rice, of which 41 are reported to be seed-borne (Ou, 1985).

Seed health testing to detect seed-borne pathogens is an important step in the management of crop diseases (Ora et al. 2011). It identifies the cause of seed infection that affects the planting value of seed lots. Farmers awareness about the effect of diseased seeds on their production is important and several routine activities need to be undertaken during storage

to reduce yield loss. These include dry seed inspection, the standard blotter test for seed infection and contamination, post entry planting for field inspection of undetected plant diseases of seed-borne and seed-contaminated pathogens (Mew and Gonzales, 2002). Seed health testing procedure involves techniques, such as direct examination of dry seeds, examination of germinated seeds, as well as examination of organism (insects and fungal). Therefore, the current survey was carried out to assess seed health status of farmers own saved seeds and to gain insight into the incidence of diseases associated with rice production in varied agro-climatic conditions of Himachal Pradesh.

2. METHODOLOGY

Roving surveys between 2022 and 2023 were undertaken in eight districts of Himachal Pradesh viz., Bilaspur, Chamba, Hamirpur, Kangra, Kullu, Mandi, Sirmour and Una. Various Rice diseases, including false smut, bacterial leaf blight, neck blast, sheath rot, sheath blight, brown spot, and leaf scald, were surveyed. The data on disease incidence and variety infected was collected by assessing upper three leaves of each random tiller from each of the ten random hills from each field and expressed as per cent for each location (Dhruw and Sahu, 2023)

$$\text{Disease Incidence (\%)} = \frac{\text{No. of diseased leaves}}{\text{Total no. of leaves assessed}} \times 100$$

The experiments were conducted at the seed pathology laboratory of the Department of Plant Pathology, Chaudhary Sarwan Kumar Himachal Pradesh Krishi Vishvavidyalaya, Palampur, India between 2022 and 2023 with seeds of nineteen exotic varieties and 1 local variety. Seed samples of exotic varieties include, HPR 2720, Sawa 200, PAC 807 plus, HPR 2612, Az 6508, Shahi Dawat, MRP 5632 (Suruchi), PAC 834, HPR 2795, US 312, US 315, Az 6444, Dhanya 834, Varsha Gold, Hyb 25P35, Hyb 257, Hyb 1067, Star 795 and LG No. 1 were obtained from farmers saved seeds to examine the health

status. Seed samples were analyzed for the detection of seed borne fungi by following the standard blotter methods by International Seed Testing Agency (ISTA) (Aremu et al. 2021). In Blotter paper method, three pieces of filter paper were soaked in sterilized distilled water and placed at the bottom of 9cm diameter sterilized Petri-dishes after draining out excess water. The experiment was set up in a completely randomized design. Nearly four hundred seed samples were analyzed for presence of seed borne pathogen(s). Seeds of each sample both sterilized (s) and unsterilized (Us), were used in this method with four replications (100 seeds per replication). Sterilization of seeds was done by immersing seeds in sodium hypochlorite solution (1%) for 3 minutes followed by three subsequent washings with distilled water. Seeds were then placed on the moist filter paper at the rate of 10 seeds per Petri dish. The Petri dishes were then incubated at ambient conditions of 25+1°C for seven days. After 7 days of incubation, the seeds were examined for fungal growth under microscope. Slides of different fungal spores encountered were prepared and examined with compound microscope at x40 magnification to establish identity of each fungus. Percentage occurrence of each fungi noticed was recorded.

3. RESULTS AND DISCUSSION

3.1 Surveys during *Kharif* 2022 and 2023

The results of the disease survey in rice producing areas of Himachal Pradesh revealed that rice seeds collected from various rice growing farmers were infected with fungi organisms which affect the germinability percentage in varying degrees. In 2022, the incidence of false smut, sheath rot was low (5-10%), while high incidence of neck blast disease (15-20%) was observed in Sawa 200 in Kangra district (Fig. 1). In Mandi district, the incidence of neck blast was low (10-15%), with a high incidence of false smut disease (20-25%) and sheath rot (15-20%) observed in US 312, US 315 in Balh. In Sirmour district, the incidence of neck blast, sheath rot, sheath blight, brown spot was low (5-10%), with a high incidence of false smut disease (upto 50%). In Una district, the incidence of sheath blight, narrow brown leaf spot was low (3-5%), with a high incidence of sheath rot (10-15%), false smut (upto 30%) observed in Hybrids 25P35, Az 6444 variety (Table 1). In Chamba district, the incidence of false smut, narrow brown leaf spot was low (5-10%), with a

high incidence of sheath rot (10-15%) observed in PAC 807 plus variety. In Bilaspur district, the incidence of neck blast was low (1-10%) observed in local grown variety. In Kullu district, the incidence of false smut was low (5-10%), with a high incidence of neck blast (10-15%) observed in local variety. In Hamirpur district, the incidence of false smut (5-30%) and (10-25%) brown spot was high in LG No. 1 and Star 795. Rice stunting has been observed as an emerging disease in almost all major rice growing areas of Himachal Pradesh affecting almost all the cultivated varieties in the farmers' fields including the hybrids. All the varieties/hybrids viz., Sawa 134, Sawa 234, Buland Raja 88, PR 121 were infected with stunting disease. No effect of date of planting was observed on the severity of stunting in most of the rice varieties. Furthermore, during year 2023, bacterial leaf blight, brown spot and false smut incidence was recorded in Sirmour and Kangra districts of Himachal Pradesh. Brown Spot, narrow brown and false smut were also prevalent in all the districts surveyed. However, the incidence of blast was negligible across all the districts with patchy occurrence to an extent of 5 per cent distribution in Kangra and Sirmour districts. Prevalence of other diseases like sheath rot, sheath blight were noticed. Incidence of leaf scald disease was observed in few pockets in surveyed areas of Kangra district in Basmati varieties. Rain, overcast weather, and high relative humidity provide favourable conditions for disease development (Magar, 2015). The intensity of biotic stresses in rice production is increasing at a startling pace because of rapid changes in climate (Jamaluddin et al. 2020). Changing climatic conditions are contributing to the emergence of new virulent pathogen strains and the occurrence of diseases in different localities. Many diseases considered as minor thus far have gained significant economic importance in many rice-growing regions, and are exacerbating their impact (Anderson et al. 2004).

3.2 Seed Health Status of Farmers own Saved Seeds in Rice

Nearly 400 samples of farmer saved seeds were analyzed for checking the seed health status. In samples collected and analyzed from Hamirpur district revealed 30% incidence of false smut and 25% incidence of brown spot across varieties such as Sawa 200, Star 795, and LG No. 1. Furthermore, samples from Kangra district exhibited 17.31% incidence of brown spot and

30.77% incidence of false smut, encompassing varieties including HD 3086, HS 562, 5258, GBW 343, 6129, Tanatan, Star 795, Sawa 200, LV, 234No., 200 No., and Shriram (Fig. 2). In Mandi district, analysis of samples showed 33.33% incidence of false smut and 29.63% incidence of brown spot among varieties such as Tanatan, Star795, Sawa 200, LV, and Chaina. Similarly, Chamba district's samples exhibited a 15.38% incidence of false smut and 7.69% incidence of

brown spot across Vijeta 700 and LV varieties. Meanwhile, Una district's samples demonstrated a 50% incidence of false smut and a 20% incidence of brown spot within PR 126 and Star 795 varieties. Lastly, in Sirmour district, examination of samples revealed a 45% incidence of false smut and a 47.5% incidence of brown spot among varieties such as Arize 6444, Pusa 1509, HPR 2795, PR 130, PR126, Lal Killa, Pusa 1121, and Sarvati.



Fig. 1. Diagnostic symptoms observed during surveys of different rice disease during *kharif* 2022 and 2023

Table 1. Status of diseases in rice hybrids/ improved rice varieties cultivated by farmers in Himachal Pradesh during *kharif* 2022 and 2023

Location District/ Blocks (Latitude & Longitude)	Location	Variety	Disease Incidence (%)					
			False Smut	Bacterial Leaf Blight (BLB)	Neck Blast	Sheath Rot	Sheath Blight	Brown Spot (BS)/ Narrow Brown Leaf Spot (NBLS)
Kangra								
Rait (32°11'03"N 76°12'47"E)	Rehlu	HPR 2720	5-10	-	-	5-10	-	-
	Shahpur	Sawa 200	5-10	-	15-20	5-10	-	-
	Shahpur	PAC 807 plus	5-10	-	-	-	-	-
	Dohab	PAC 807 plus	5-10	-	-	-	-	-
Dharamshala (32°13'00"N 76°19'08"E)	Bandi	Sawa 200, PAC 807 plus	10-15	-	5-10	-	-	-
	Gharoh	HPR 2612	-	-	5-10	-	-	-
		HPR 2720	5-10	-	-	5	-	-
	Jadrangal	Local, hybrid	10-15	-	-	5-10	-	-
	Banoi	Az 6508	10-15	-	-	-	-	-
Nagrota Bagwan (32°06'29"N 76°22'56"E)	Chahri	Sawa 200, Star 795	10-15	-	40-50	10-15	-	-
	Malan	Az 6508	10-15	-	-	-	-	-
	Massal	Sawa 200, Shahi Dawat	10-15	-	25-30	5-10	-	-
	Pathiar	MRP 5632 (Suruchi)	10-15	-	-	5-10	-	-
Bhawarna (32°02'41"N 76°29'33"E)	Chimbalhaar	PAC 807 plus	<5	-	-	-	-	-
		HPR 2612	-	-	-	-	-	-
	Bagora	Sawa 200, LG No.1	5-10	-	5-10	5-10	-	-
		PAC 834	5-10	-	-	-	-	-
	Mainjha	PAC 834, Sawa 200	5-10	-	-	5-10	-	-
Mandi								
Sandhol (31°51'31"N 76°39'09"E)	Sandhol	HPR 2795	-	-	-	-	-	-
	Lasrana	Hybrid (Not known)	15-20	-	10-15	15-20	-	15-20 (BS); 40-50 (NBLS)
Balh (31°53'22"N 76°55'15"E)	Galma	US 312	25-30	-	-	5-10	-	10-15 (BS)
	Sundernagar	US 312, US 315	25-30	-	-	5-10	-	-
Sirmour								
Paonta Sahib	Dhaulakuan	Az 6444	25-30	-	5-10	5-10	10-15	10-15 BS

Location District/ Blocks (Latitude & Longitude)	Location	Variety	Disease Incidence (%)					
			False Smut	Bacterial Leaf Blight (BLB)	Neck Blast	Sheath Rot	Sheath Blight	Brown Spot (BS)/ Narrow Brown Leaf Spot (NBLS)
(30°31'54"N 77°34'02"E)	Puruwala	Az 6444, Dhanya 834	25-30	-	5-10	5-10	5-10	10-15
	Surajpur	Az 6444, Varsha Gold	30-35	-	5-10	5-10	10-15	5-10
	Rampur	Hyb 25P35	Up to 50%	-	5	10-15	5-10	5-10
	Kolar	Az 6444, Dhanya 834, Varsha Gold	25-30	-	5-10	5-10	10-15	15-20
Una								
Una (31°28'14"N 76°16'07"E)	Rampur	Az 6444	10-15	-	-	-	-	5-10 NBLS/ BS
	Basal	Hyb. 25P35, Az 6444	10-20	-	-	-	-	10 BS
	Nangal Khurd	Hyb. 257, Hyb. 1067	25-30	-	-	-	-	10-15 NBLS/ BS
	Nandpur	Az 6444	7-10	-	3	12-15	3	5
	Ispur	Az 6444	3	-	3	12	3	5
	Andora	Hyb. 25P35, Az 6444	8	-	5	10	5	5
	Kuthiari	Hyb. 257, Hyb. 1067	10	-	3	10	5	4
	Chak	Az 6444	5	-	4	10	4	4
Chamba	Thathal	Az 6444	5	-	4	8	4	4
	Bhatiyat (32°26'57"N 76°00'37"E)	PAC 807 plus	5-10	-	-	10-15	-	5-10 NBLS
Bilaspur								
Bilaspur (22°04'46"N 82°08'27"E)	Jhandutta	Local	-	-	5-7	-	-	-
	Bilaspur sadar	Local	-	-	2-3	-	-	-
	Ghumarwin	Local	-	-	10-11	-	-	-
	Nainadevi	Local	-	-	1-3	-	-	-
Kullu								
Kullu (31°57'16"N 77°06'40"E)	Baragran	Local	10	-	15	-	-	-
	Chhaki	Local	5	-	10	-	-	-
	Naggar Seri	Local	10	-	15	-	-	-
	Haripur	Local	10	-	15	-	-	-

Location District/ Blocks (Latitude & Longitude)	Location	Variety	Disease Incidence (%)					
			False Smut	Bacterial Leaf Blight (BLB)	Neck Blast	Sheath Rot	Sheath Blight	Brown Spot (BS)/ Narrow Brown Leaf Spot (NBLS)
Hamirpur								
Hamirpur (31°68'62"N 76°52'13"E)	Bhira	Sawa 200	10-15	-	-	-	-	10-15
	Kehdru	Local	25-30	-	-	-	-	10-15
Bhoranj (31°64'30"N 76°64'27"E)	Khad Bazar	Star 795	7-10	-	-	-	-	20-25
	Dimmi	LG No. 1	25-30	-	-	-	-	5-10
	Bhareri	Local	10-15	-	-	-	-	10-15



Fig. 2. Microscopic observation of mycoflora associated with false smut, blast and brown spot from farmers saved rice seeds

Seed-borne infections in rice cause 50-80% yield losses, depending on the agroecology, disease severity, and crop susceptibility (Monajjem et al. 2014). Farmers who rely on their farm-saved seeds are likely to contaminate their subsequent seeds for the next planting season in instances where there is seed-borne pathogen transmission from the soil. Most seed-borne diseases caused by the fungi are disastrous as they may decrease seed germination, causing seed discoloration and produce toxins that may be detrimental to man and domestic animals. Association of mycoflora i.e., *Ustilaginoidea virens*, *Pyricularia oryzae* and *Helminthosporium oryzae* in farmer saved seed was also observed (Fig. 2). Distribution of some seed-borne fungi associated with rice seeds has been reported in many countries, including Nigeria (Suleiman & Omafè, 2013); Pakistan (Butt et al. 2011); Egypt (Madbouly et al. 2012); Bangladesh (Ora et al. 2011); Cameroon (Nguefack et al. 2007) and Chad Republic (Serferbe et al. 2016). The results are in agreement with that of Utobo et al. (2011) who reported that *T. padwickii*, *Ustilaginoidea virens* and *Helminthosporium oryzae* were the most frequently isolated seed borne fungi irrespective of the source of the rice variety tested in their study. Singh & Brar (2020) recorded minimum neck blast incidence (14%) in the variety PR 126. Aishwarya et al. (2023) detected and characterized different pathogenic isolates of brown spot and blast disease in Tamil Nadu.

Major resistance genes from different resistance donors have been reported for various rice diseases. To date, over 44 resistance genes have been identified against bacterial leaf blight (Kim and Reinke, 2019). More than 100 distinct blast-resistance genes have been reported across different rice chromosomes, of which 21 have been cloned (Devi et al. 2020). However,

no single resistance gene confers resistance to sheath blight. Two major sheath blight QTLs (qShB9-2 and qSBR11-1) have been reported to contribute more than 10% to sheath blight resistance (Jamaloddin et al. 2021). But, thus far, genetic diversity for high resistance/tolerance of sheath blight has not been reported in either cultivated rice or its wild relatives (Bonman, 1992). For bacterial leaf streak (BLS), no major resistance genes (R-genes) have been identified (Bossa-Castro et al. 2018). These resistance genes, though identified in different studies, play a highly connected role in enabling rice to combat invading pathogens. Consequently, the presence or absence of resistance genes significantly influences disease incidence in rice plants infected by various pathogens.

4. CONCLUSION

Roving survey and laboratory experiments were conducted between 2023 and 2024 to determine the occurrence of seed borne-fungi over both the year under study. Standard blotter method was used for assessment. Fungi detected were *Ustilaginoidea virens*, *Pyricularia oryzae* and *Helminthosporium oryzae*. It is evident from the study that seed-borne fungi were associated with all the rice cultivars sampled from the study area. There was high prevalence of storage fungi associated with the rice seeds partly because of the storage practices employed by the farmers. The need to emphasize the role of healthy seed in establishment of a healthy crop is highlighted with this work.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image

generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENT

The first author is thankful to Indian Council of Agricultural Research (ICAR) for providing financial support to undertake this study through "AICRP on Seed (Crops)" (ICAR-017-17).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Aishwarya, V., Manoranjitham, S. K., Gopalakrishnan, C., Somasundaram, E., & Geethanjali, S. (2023). Isolation and characterization of rice blast and brown spot disease in different regions of Tamil Nadu, India. *International Journal of Plant & Soil Science*, 35(19), 975-985.
- Akter, M., & Hossain, I. (2015). Quality of some hybrid seeds of rice and control of seed-borne fungi in Bangladesh. *Journal of the Bangladesh Agricultural University*, 13(2), 161-168.
<https://doi.org/10.3329/jbau.v13i2.28771>
- Anderson, P. K., Cunningham, A. A., Patel, N. G., Morales, F. J., Epstein, P. R., & Daszak, P. (2004). Emerging infectious diseases of plants: Pathogen pollution, climate change and agrotechnology drivers. *Trends in Ecology & Evolution*, 19, 535-544.
- Anselme, C. (1981). The importance in cultivation of pathogenic organisms transmitted by seeds. *Seed Science and Technology (Netherlands)*, 9, 689-695.
- Aremu, M. B., Kurah, I. A., Okparavero, N. F., Okunade, S. O., Adebola, M. O., Rahman, M. O., & Aina, O. B. (2021). Assessment of seed-borne plant pathogenic fungi, insect emergence and percentage germinability associated with harvested and stored rice in Kwara State, Nigeria. *African Journal of Agriculture and Food Science*, 4(2), 44-54.
<http://doi.org/10.52589/AJAFS-OY9AE3MW>
- Bonman, J. M. (1992). Durable resistance to rice blast disease-environmental influences. *Euphytica*, 63, 115-123.
<https://doi.org/10.1007/BF00023917>
- Bossa-Castro, A. M., Tekete, C., Raghavan, C., Delorean, E. E., Dereeper, A., Dagno, K., Koita, O., Mosquera, G., et al. (2018). Allelic variation for broad-spectrum resistance and susceptibility to bacterial pathogens identified in a rice MAGIC population. *Plant Biotechnology Journal*, 16, 1559-1568.
<https://doi.org/10.1111/pbi.12895>
- Butt, A., Yaseen, S., & Javaid, A. (2011). Seed-borne mycoflora of stored rice grains and its chemical control. *Journal of Animal and Plant Sciences*, 21(2), 193-196.
- Dede, Y. K., Ahmad, J., Iskandar, L., & Titi, C. S. (2019). Evaluation of growth and physiological responses of three rice (*Oryza sativa* L.) varieties to elevated temperatures. *Journal of Tropical Crop Science*, 6(1), 17-23.
<https://doi.org/10.29244/jtcs.6.01.17-23>
- Devi, S. J. S. R., Singh, K., Umakanth, B., Vishalakshmi, B., Rao, K. V. S., Suneel, B., Sharma, S. K., et al. (2020). Identification and characterization of a large effect QTL from *Oryza glumaepatula* revealed Pi68(t) as putative candidate gene for rice blast resistance. *Rice*, 13, 17.
<https://doi.org/10.1186/s12284-020-00378-4>
- Dhruw, S., & Sahu, S. (2023). Assessment of brown spot disease incidence and severity in rice cultivation across different regions of Chattisgarh. *The Pharma Innovation Journal*, 12(11), 1663-1667.
- Islam, M. S., Jahan, Q. S. A., Bunnarith, K., Viangkum, S., & Merca, S. D. (2000). Evaluation of seed health of some rice varieties under different conditions. *Botanical Bulletin of Academia Sinica*, 41, 293-297.
- Jamaloddin, M., Durga, C. V., Swathi, G., Anuradha, C., Vanisri, S., Rajan, C. P. D., Raju, S. K., et al. (2020). Marker assisted gene pyramiding (MAGP) for bacterial blight and blast resistance into mega rice variety "Tellahamsa". *PLoS One*, 15, e0234088.
<https://doi.org/10.1371/journal.pone.0234088>
- Jamaloddin, M., Mahender, A., Gokulan C. G., Balachiranjeevi, C., Maliha, A., Patel, H. K., & Ali, J. (2021). Molecular approaches for disease resistance in rice. In J. Ali & S. H. Wani (Eds.), *Rice Improvement* (pp. 215-250). Springer.
https://doi.org/10.1007/978-3-030-66530-2_10
- Kim, S. M., & Reinke, R. F. (2019). A novel resistance gene for bacterial blight in rice,

- Xa43(t) identified by GWAS, confirmed by QTL mapping using a bi-parental population. *PLoS One*, 14, e0211775. <https://doi.org/10.1371/journal.pone.0211775>
- Madbouly, A. K., Ibrahim, M. I., Sehab, A. F., & Abdel-Wahhab, M. A. (2012). Co-occurrence of mycoflora, aflatoxins, and fumonisins in maize and rice seeds from markets of different districts in Cairo, Egypt. *Food Additives and Contaminants*, 5(2), 112-120. <https://doi.org/10.1080/19393210.2012.676078>
- Magar, P. B. (2015). Screening of rice varieties against brown leaf spot disease at Jyotinagar, Chitwan, Nepal. *International Journal of Applied Sciences and Biotechnology*, 3(1), 56-60.
- McGee, D. C. (1995). Epidemiological approach to disease management through seed technology. *Annual Review of Phytopathology*, 33, 445-466.
- Mew, T. W., & Gonzales, P. (2002). A handbook of rice seed borne fungi. International Rice Research Institute, Los Baños, Philippines, 83.
- Monajjem, S., Zainali, E., Ghaderi-Far, F., Soltani, E., Chaleshtari, M. H., & Khoshkdam, M. (2014). Evaluation seed-born fungi of rice (*Oryza sativa* L.) and their effect on seed quality. *Journal of Plant Pathology & Microbiology*, 5(4), 1-7. <https://doi.org/10.4172/2157-7471.1000239>
- Nguefack, J., Nguikwie, S., Fotio, D., Dongmo, B., Zollo, P. A., Leth, V., et al. (2007). Fungicidal potential of essential oils and fractions from *Cymbopogon citratus*, *Ocimum gratissimum*, and *Thymus vulgaris* to control *Alternaria padwickii* and *Bipolaris oryzae*, two seed-borne fungi of rice (*Oryza sativa* L.). *Journal of Essential Oil Research*, 19(6), 581-587. <https://doi.org/10.1080/10412905.2007.9699336>
- Ora, N., Faruq, A. N., Islam, M. T., Akhtar, N., & Rahman, M. M. (2011). Detection and identification of seed-borne pathogens from some cultivated hybrid rice varieties in Bangladesh. *Middle-East Journal of Scientific Research*, 10(4), 482-488. <https://hdl.handle.net/10536/DRO/DU:30102133>
- Ou, S. H. (1985). *Rice diseases* (2nd ed.). Kew, England: Commonwealth Mycological Institute, 368-380.
- Serferbe, S., Tsopmbeng, N. G., & Kuate, J. R. (2016). Seed-borne fungi associated with rice seeds varieties in Bongor, Chad Republic. *International Journal of Current Microbiology and Applied Sciences*, 5(12), 161-170. <http://dx.doi.org/10.20546/ijcmas.2016.512.018>
- Singh, N., & Brar, J. S. (2020). Screening of disease incidence in rice cultivars against *Pyricularia oryzae* in Talwandi Sabo, Punjab, India. *International Journal of Current Microbiology and Applied Sciences*, 9(08), 2198-2208. <https://doi.org/10.20546/ijcmas.2020.908.251>
- Suleiman, M. N., & Omafefe, O. M. (2013). Activity of three medicinal plants on fungi isolated from stored maize seeds (*Zea mays* L.). *Global Journal of Medicinal Plant Research*, 1, 77-81.
- Udemezue, J. C. (2018). Analysis of rice production and consumption trends in Nigeria. *Journal of Plant Science and Crop Protection*, 1(3), 305.
- USDA. (2024). *World Agricultural Production*.
- Utobo, E. B., Ogbodo, E. B., & Nwogbaga, A. C. (2011). Seed borne mycoflora associated with rice and their influence on growth at Abakiliki, Southeast agro-ecology, Nigeria. *Libyan Agriculture Research Centre – Journal International*, 2(2), 79-84.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/128744>