



# **Sustainable Technologies for Soil Health and Basmati Rice Productivity in India: Current Research and Future Directions**

**Shivangi <sup>a++\*</sup>, Omkar Singh <sup>b#</sup>, UP Shahi <sup>b†</sup>, P K Singh <sup>c‡</sup>,  
KK Singh <sup>a^</sup>, Pintoo Kumar <sup>a##</sup>, Shweta Singh <sup>b#^</sup>,  
Vaishali Singh <sup>b§</sup>, Adarsh Kumar Meena <sup>b§</sup>  
and Akansha Tyagi <sup>b§</sup>**

<sup>a</sup> KVK Bijnor, India.

<sup>b</sup> Department of Soil Science, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut, U.P., India.

<sup>c</sup> Directorate of Extension, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut, U.P., India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/acri/2025/v25i51208>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/135220>

<sup>++</sup> SMS (Agronomy);

<sup>#</sup> Senior Research Fellow;

<sup>†</sup> Professor & Head;

<sup>‡</sup> Director Extension;

<sup>^</sup> SMS (Plant Breeding);

<sup>##</sup> SMS (Plant Protection)

<sup>#^</sup> Research Associate;

<sup>§</sup> Research Scholar;

\*Corresponding author: Email: [Singhomkar.agri@gmail.com](mailto:Singhomkar.agri@gmail.com); [singhshivangi.agri@gmail.com](mailto:singhshivangi.agri@gmail.com);

**Cite as:** Shivangi, Omkar Singh, UP Shahi, P K Singh, KK Singh, Pintoo Kumar, Shweta Singh, Vaishali Singh, Adarsh Kumar Meena, and Akansha Tyagi. 2025. "Sustainable Technologies for Soil Health and Basmati Rice Productivity in India: Current Research and Future Directions". Archives of Current Research International 25 (5):274-95. <https://doi.org/10.9734/acri/2025/v25i51208>.

## Review Article

Received: 29/02/2025  
Accepted: 02/05/2025  
Published: 05/05/2025

## ABSTRACT

Basmati rice (*Oryza sativa* L.) is a crucial economic and cultural crop in South Asia, particularly in India and Pakistan. But the intensification of basmati rice farming methods has resulted in notable soil degradation, lower soil organic carbon, imbalances of nutrients, and less biodiversity. The present situation of sustainable technologies meant to improve soil condition and increase basmati rice production is investigated in this thorough research. Research on conservation agriculture, organic amendments, microbial inoculants, precise nutrient management, and climate-smart agricultural methods is compiled in this work. By means of field investigations, meta-analyses, and case studies, we have found interesting approaches concurrently addressing soil health improvement and yield optimisation. This review emphasizes important information gaps including limited long-term studies in many agroecological areas, insufficient economic analysis, and poor evaluation of technological adoption hurdles. Emphasizing the requirement of integrated approaches combining several sustainable technologies, increased stakeholder involvement, and policy frameworks encouraging sustainable soil management practices should be the goal of future studies. This thorough study offers evidence-based recommendations for creating all-encompassing plans for sustainable basmati rice output systems to researchers, legislators, and industry experts.

**Keywords:** *Basmati rice; soil health; conservation agriculture; organic amendments; microbial inoculants; precision agriculture; sustainable intensification.*

## ABBREVIATIONS

ZT	: Zero Tillage
RR	: Residue Retention
SSNM	: Site-Specific Nutrient Management
AWD	: Alternate Wetting and Drying
DSR	: Direct-Seeded Rice
GM	: Green Manure
PGPR	: Plant Growth-Promoting Rhizobacteria
NI	: Nitrification Inhibitors
CA	: Conservation Agriculture
FYM	: Farmyard Manure
BI	: Biofertilizer Inoculation
VC	: Vermicompost
BC	: Biochar
SOC	: Soil Organic Carbon
MBC	: Microbial Biomass Carbon
WSA	: Water Stable Aggregates
BD	: Bulk Density
GHG	: Greenhouse Gas Emissions

## 1. BACKGROUND

Basmati rice (*Oryza sativa* L.) is one of the most distinguished aromatic rice varieties globally, and is characterized by its distinctive fragrance, flavor profile, and remarkable grain elongation upon cooking (Bhattacharjee et al., 2022; Singh et al. 2023). Mostly grown on the Indo-Gangetic Plains

between northern India and Pakistan, basmati rice is a major export rice speciality from around the world and commands premium pricing in foreign markets (Dar et al., 2020). With a projected market value of US\$6.8 billion, global basmati exports in 2023 topped 6.4 million metric tonnes, therefore highlighting its economic importance in the agricultural export portfolios of producing countries (Singh O., 2023; FAO, 2024; Singh et al., 2024). Beyond its economic worth, basmati rice has great cultural and historical importance in South Asia, shown mostly in traditional cuisines, religious events, and cultural festivities (Singh et al., 2018; Singh et al., 2023). Basmati rice's geographical indication (GI) status given in 2016 acknowledges its special connection with particular areas of the Indo-Gangetic Plains, where unique soil features, climatic conditions, and traditional farming methods help to define exceptional quality criteria (Giraud, 2013; Singh et al., 2018; Dutta et al., 2024).

With the Indian states of Punjab, Haryana, and Uttar Pradesh alongside Pakistan's Punjab province, roughly 90% of the world's basmati rice is produced in the north-west of the Indian subcontinent (Mahajan et al., 2021; Singh et al., 2023; Singh et al., 2024). These areas'

agroecological conditions—alluvial soils, different temperature variances between day and night, and suitable water availability—create an ideal habitat for developing the unique grain quality and aromatic compounds that define basmati rice (Singh et al., 2018; Singh et al., 2024; Singh O. 2023).

In modern agricultural systems, basmati rice output presents major sustainability issues notwithstanding their economic and cultural value (Balafoutis et al., 2020). While significantly raising production volumes, the intensification of rice farming in the era of the green revolution has resulted in major ecological effects over basmati-growing areas (Ladha et al., 2016; Bünemann et al., 2018). Particularly puddle operations involving repetitive ploughing of flooded fields, intensive tillage methods have drastically changed the soil structure and hydrological parameters (Gathala et al., 2011; Mugabo et al., 2024; Jagadish et al., 2015). Compacted plough pans, lower macroporosity, and less populations of beneficial soil organisms have emerged from this (Sharma et al., 2019; Singh et al., 2019; Ghazaryan et al., 2024). Mostly seen in basmati-growing areas, constant rice-wheat farming methods have resulted in extreme nutrient imbalances in agricultural soils. With average contents of 0.37–0.52%, much below the threshold (0.75%), long-term fertiliser experiments carried out over several sites in the Indo-Gangetic Plains show trends in declining soil organic carbon (SOC) levels (Benbi and Brar, 2009; Bhattacharyya et al., 2015; Lipper et al., 2014). With a utilisation efficiency sometimes below 35%, nitrogen application in conventional basmati farming often surpasses 180 kg N/ha and causes significant nutrient losses to the environment through leaching, volatilisation, and denitrification processes (Timsina et al., 2016; Rodell et al., 2009). Gupta et al. (2019) conducted studies on soil acidification in intensively grown basmati fields, with a pH drop of 0.5–1.2 units noted over a decade in the provinces of Punjab and Haryana.

Climate change adds still another level of complication to basmati rice farming operations. Productivity and sustainability are seriously threatened by expected rises in temperature, changed precipitation patterns, and more frequency of extreme weather events (Kumar et al., 2020; Sachan et al., 2023; Singh et al., 2023; Sanjeev et al., 2020). Whereas higher nighttime temperatures negatively affect grain quality

metrics, including fragrance compounds and amylose concentration, heat stress during blooming can diminish spikelet fertility by 35–45%. Climate models show that, without adaptation strategies, impacts of climate change might cause basmati yields to drop by 15–30% by 2050 over most growing areas (Aryal et al., 2020). Basmati rice farming's financial considerations provide still another difficulty (Subash et al., 2023). Rural-urban migration and rivalry from non-agricultural industries have caused labour expenses to rise by 150–200% over the past ten years, so sharply escalating production costs (Mahajan et al., 2021; Shivangi et al., 2025a). Particularly with regard to transplanting and harvesting, the availability of agricultural labour during important operations has become increasingly limited, which calls for mechanization and other establishment techniques (Jat et al., 2020). Further complicating the economic environment of basmati growers include market volatility, worldwide competitiveness, and changing customer tastes for sustainability certifications (Singh et al., 2018).

Rising as a major paradigm in sustainable agriculture, the idea of soil health offers a comprehensive approach to soil management transcending fertility concerns to include the capacity of soil to function as a vital living ecosystem (Doran and Zeiss, 2000; Shivangi et al., 2025b). From this point of view, soil is a dynamic, complex system whose capacity depends on the combined performance of physical, chemical, and biological characteristics supporting several ecosystem services (Kibblewhite et al., 2008; Karlen et al., 2016). In agroecosystems, soil health refers to several vital roles: sustaining plant productivity through nutrient cycling and structural support; controlling water infiltration and storage; sequesters carbon; reduces greenhouse gas emissions; filters pollutants; provides habitat for biodiversity (Lehmann et al., 2020). Soil health indices shown very high relationships with productivity and sustainability measures in basmati rice growing systems. With each 0.1% rise in SOC related with yield gains of 0.2–0.3 t/ha, Sharma et al. (2021) found strong positive correlations between SOC content and grain yield. Analogously, Yadav et al. (2019) showed that farms with balanced nutrient profiles yielded 15–25% more than those with several nutritional deficits. With microbial biomass carbon explaining 53–67% of yield changes across various management methods, the biological

aspects of soil health exhibit similarly important correlations with production (Rana et al., 2020). The multifarious character of healthy soils offers chances for cooperative methods towards agricultural sustainability. Many times, practices that improve soil health also help climate resilience, resource-use efficiency, biodiversity, and environmental footprint (Lal, 2020). This possibility to simultaneously solve several sustainability issues has made soil health a central focus for policies, research projects, and agricultural growth plans all around (Lehmann et al., 2020).

With many conceptual frameworks directing research and development efforts, sustainable rice farming paradigms have seen notable evolution in recent years. Pretty and Bharucha's (2014) sustainable intensification strategy stresses the concurrent improvement of production and environmental outcomes via knowledge-intensive approaches rather than higher input consumption. For basmati rice systems, where premium quality criteria call for methods that improve both production and qualitative criteria while lowering environmental impacts, this concept is especially pertinent (Pretty et al., 2018). Described by Altieri et al. (2015), the agroecological paradigm stresses on biodiversity, synergy, resource efficiency, and resilience by means of redesign of farming systems based on ecological principles. Applied to basmati rice, this method stresses diversification by rotation, integration of legumes, habitat management for beneficial creatures, and use of local knowledge systems (Singh et al., 2020). With methods meant to rebuild soil organic matter, restore degraded soil biodiversity, and improve ecosystem functioning, the regenerative agriculture movement extends this perspective to specifically emphasise soil health restoration as a fundamental objective (LaCanne and Lundgren, 2018).

From agricultural methods to biological interventions to technical applications to knowledge systems, the shift towards sustainable technologies in basmati rice systems spans a broad spectrum of developments. Zero or less tillage, direct sowing, and residue retention among conservation agriculture techniques have shown promise to solve soil physical degradation and improve resource use efficiency (Jat et al., 2020; Singh et al., 2023). Field trials over the Indo-Gangetic Plains show that, following an initial transition period, the implementation of conservation agriculture

practices can lower production costs by 15–25% while either maintaining or improving yields (Gathala et al., 2021). Organic additions comprising farmyard manure, compost, vermicompost, and green manure represent established, but changing ways to improve soil biological functioning and nutrient cycle (Bhattacharyya et al., 2021; Shivangi et al., 2023; Saikant et al., 2023). Along with notable improvements in soil health metrics (Singh et al., 2021; Singh et al., 2022; Rai et al., 2023), the combination of these amendments with lowered inorganic fertilizer applications has shown yield increases of 8–15% compared to mono inorganic fertilization. Microbial inoculants provide biological means to improve nutrient availability and plant resilience: nitrogen-fixing bacteria, phosphate-solubilizing microorganisms, arbuscular mycorrhizal fungus, and plant growth-promoting rhizobacteria (Gouda et al., 2018). With field evaluations showing yield increases of 10–20% when used in optimal combinations, these biologicals are increasingly being structured as consortia addressing several functions simultaneously (Rai et al., 2020). Particularly alternate wetting and drying (AWD), subsurface drip irrigation, and laser land levelling, water management advances directly address important water sustainability concerns in basmati production (Jain et al., 2019). Comparatively to conventional flooding methods, these methods have shown water savings of 20–35% while keeping yields and lowering greenhouse gas emissions (Jat et al., 2020).

Digital and precise technologies—including GPS-guided machinery, drone-based surveillance, sensor networks, and decision support systems—increasingly enable knowledge-intensive management approaches that maximise resource usage according to site-specific needs (Balyan et al., 2019). Site-specific nutrient management (SSNM), which has shown yield increases of 10–15% and nitrogen use efficiency by 25–40% compared to standard techniques (Pampolino et al., 2019), is easier to execute with these technologies. With evidence pointing to suitable combinations creating synergistic advantages that surpass the total of individual therapies, the integration of several technologies into coherent systems is a particularly exciting future (Jat et al., 2020). Nevertheless, the context-specificity of ideal technology packages calls for flexible solutions fit to local agroecological conditions, resource availability, market access, and farmer capability (Sharma et al., 2021; Shahi et al., 2024).

Synthesizing present research on methods that simultaneously improve soil health and preserve or increase production and quality is desperately needed given the economic value of basmati rice and the difficult sustainability issues related with its farming. This thorough evaluation seeks to assess the efficiency of several sustainable technologies, point up knowledge gaps, and suggest future study paths to influence the creation of overall strategies for sustainable basmati rice-production systems. This review analyses the current state of soil health in basmati rice cultivation systems, identifying key constraints and their implications for long-term sustainability; evaluates the effectiveness of sustainable technologies for enhancing soil health and basmati rice productivity, including conservation agriculture, organic amendments, microbial inoculants, precision nutrient management, and climate-smart agricultural practices; assesses the potential for integrating multiple technologies to maximize agronomic, economic, and environmental benefits; identifies knowledge gaps and limitations in existing research that constrain the development and adoption of sustainable technologies; and proposes future research directions and policy recommendations for promoting sustainable soil management in basmati rice production. This review helps to further the more general aim of balancing productivity objectives with environmental sustainability and climate resilience in premium rice production by critically analyzing the present knowledge and pointing up interesting avenues for future research and development.

## **2. PRESENT STATE OF SOIL HEALTH IN SYSTEMS OF BASMATI RICE PRODUCTION**

### **2.1 Physical Indicators of Soil Health**

The physical characteristics of the soil have been greatly changed by intensive farming methods used in classic basmati rice systems. Numerous studies have found declining soil structure; Singh et al. (2019) found higher bulk density (1.45–1.62 g/cm<sup>3</sup>) in intensive rice-wheat systems than in less intensive systems (1.32–1.41 g/cm<sup>3</sup>). Common in basmati rice farming, puddling results in a compacted plough layer that limits root penetration and water absorption (Gathala et al., 2011). Kumar et al. (2018) conducted a thorough analysis over 150 basmati rice fields in Haryana, India, finding that 67% showed poor soil aggregation and lowered

macroporosity, therefore directly influencing water penetration rates and erosion susceptibility.

### **2.2 Chemical Indicators of Soil Health**

Thus, in the basmati rice growing regions, soil chemical characteristics have been quite changed. With average values of 0.37–0.52%, long-term fertilizer studies in Punjab, India, revealed dropping soil organic carbon (SOC) levels, well below the threshold (0.75%) thought ideal for sustainable development (Benbi and Brar, 2009). With Timsina et al. (2013) reporting excessive nitrogen application (average 174 kg N/ha) combined with deficits in K, sulphur, and micronutrients across basmati-growing areas, nutrient imbalances have become somewhat common. With a pH drop of 0.5–1.2 units recorded over a decade in some areas, soil acidification has become a worry in those areas with continuous fertilizer usage (Gupta et al., 2019).

### **2.3 Biological Indicators of Soil Health**

Intensive farming methods have especially an impact on the biological elements of the soil. Just 40–60% of the amounts observed in natural or less disturbed soils are represented by microbial biomass carbon in standard basmati rice fields (187–245 µg/g). Dehydrogenase, β-glucosidase, and phosphatase are among the enzymatic activities vital for nutrient cycling that are much less in intensively managed fields (Sharma et al., 2020). Conventional fields support just 30–50% of the species richness found in organically managed systems, hence biodiversity indices for soil fauna—especially earthworms and arthropods—show a significant drop (Bhushan et al., 2019).

### **2.4 Relationships Between Soil Health and Basmati Rice Productivity**

Literature has progressively recorded the links between basmati rice production and markers of soil health. With each 0.1% rise in SOC linked with yield gains of 0.2–0.3 t/ha, Sharma et al. (2021) found robustly positive correlations between SOC content and grain yield ( $r = 0.78$ ). Analogously, Yadav et al. (2019) showed that farms with balanced nutrient profiles yielded 15–25% more than those with several nutritional deficits. With microbial biomass carbon accounting 53–67% of yield variances across

various management regimes, biological markers also demonstrate noteworthy links with productivity (Rana et al., 2020). These results highlight the need of preserving soil condition for the generation of sustainable basmati rice.

### 3. CONSERVATION AGRICULTURE FOR BASMATI RICE SYSTEMS

**Three basic ideas define conservation agriculture (CA):** Minimum soil disturbance, permanent soil cover, and crop variety (Kassam et al., 2019). CA methods in basmati rice systems include zero or less tillage, direct sowing, residue retention, and rotation-based legume or other crop inclusion (Jat et al., 2020; Dutta et al., 2023). These techniques deviate greatly from traditional approaches include puddling, intense tillage, and crop residue burning or removal.

#### 3.1 Effects on Physical Characteristics of Soil

Adoption of CA techniques has resulted in considerable changes in the soil physical state. Under zero-tillage with residue retention, Singh et al. (2021) conducted five-year research on basmati-growing areas of Haryana, India, noting lowered bulk density (from 1.58 to 1.42 g/cm<sup>3</sup>) and elevated water stable aggregates (from 38.5% to 62.3%). Comparatively to conventional tillage, the water-holding capacity rose by 12–18% and the infiltration rates improved by 45–70% (Parihar et al., 2018). < In the 15–30 cm soil layer following four years of low tillage operations, soil penetration resistance dropped by 26–35%.

#### 3.2 Affect on Chemical Properties of Soil

The chemical markers of the soil benefit from conservation agriculture. Zero tillage with residue retention raised SOC by an average of 0.27–0.39% in the top 15 cm soil layer over 3–7 years, according a meta-analysis of 26 studies in rice-based systems by Das et al. (2020) vs traditional techniques. With 15–22% greater nitrogen mineralization rates and 18–30% more phosphorus availability, nutrient availability improved greatly (Jat et al., 2019). In CA systems, soil pH stabilization has been noted under lower acidification rates than in more conventional methods (Choudhary et al., 2018).

#### 3.3 Effects on Biological Characteristics of Soil

Particularly strong responses to CA methods were shown by the biological elements of soil quality. Zero-tillage systems support 30–80% more enzyme activities than conventional tillage and 40–120% more microbial biomass carbon (Sharma et al., 2020; Singh et al., 2023). With average densities of 45–78 individuals/m<sup>2</sup> compared with 12–25 individuals/m<sup>2</sup> under conventional techniques, earthworm populations rose three- to five-fold under decreased tillage with residue retention (Kumar et al., 2018). Assessed by community-level physiological profiling, functional diversity of soil microorganisms revealed notable increase under CA techniques; Shannon diversity indices changed from 2.1–2.4 to 2.8–3.2 after five years (Bhushan et al., 2019).

#### 3.4 Impacts on Productivity and Profitability of Basmati Rice

Studies of how CA affects basmati rice production have produced mixed but typically favorable results. Direct-seeded rice under zero tillage initially lowered yields by 5–8% in the first 1–2 years but matched or exceeded conventional yields by 3–7% during the transition phase, according to a thorough study of 32 on-farm trials conducted throughout the Indo-Gangetic Plains by Jat et al. (2020). Due mostly to lower labour, fuel, and machinery needs, water productivity has grown by 25–40% while production costs have dropped by 15–25% (Gathala et al., 2021). Under CA systems, profitability rose US\$100–250/ha, a 12–30% increase above standard methods (Kakraliya et al., 2018).

#### 3.5 Challenges and Limitations

Even with its advantages, CA application in basmati rice systems presents certain difficulties. Particularly in the transition phase, which usually calls for higher herbicide use or integrated weed management techniques, weed control is still a major problem (Singh et al., 2021). Because of their high starting prices and restricted availability, specialized equipment—including zero-till drills and residue management machinery—may not be available to smallholder farmers (Jat et al., 2020). Furthermore, site-specific modifications are required since local farming practices, soil type, and climate affect

performance (Gathala et al., 2019).4. Organic Amendments for Enhancing Soil Health.

### 3.6 Types and Characteristics of Organic Amendments

Various organic amendments have been evaluated for basmati rice systems, including farmyard manure (FYM), compost, vermicompost, green manure, and crop residues. These materials differ substantially in their nutrient content, decomposition rates, and effects on soil properties. Table 1 summarizes the characteristics of common organic amendments used in basmati rice cultivation.

### 3.8 Effect on Physical Properties of Soil

Organic additions greatly enhance the physical state of the soil in basmati rice systems. Compared with inorganic fertilizer alone, FYM application at 10 t/ha/year lowered bulk density by 8-15% and raised total porosity by 10-18% according a meta-analysis of 45 research by Bhattacharyya et al. (2021). Regular organic amendment treatments resulted in water-stable aggregates rising by 30–45%. Incorporation of green manure enhanced soil structure; aggregate stability rose by 25–35% after *Sesbania aculeata* cultivation before rice transplantation (Kumar et al., 2019).

### 3.9 Effects on Chemical Properties of Soil

Organic amendments greatly improved markers of soil chemical condition. Long-term (>10 year) studies showed that compared to solitary inorganic fertilization, the combined use of FYM (10 t/ha) with lowered inorganic fertilisers raised SOC by 45–60%. Regular organic amendment treatments increase the cation exchange capacity by 15–25%, hence improving nutrient retention capacity (Gupta et al., 2019). Organic inputs have been demonstrated to buffer against acidification in intensively grown soils, hence stabilizing impacts of soil pH (Sharma et al., 2019).

### 3.10 Effects on Biological Characteristics of Soil

Applications of organic amendments clearly improved the biological indices. Under integrated

organic and inorganic nutrient management, microbial biomass carbon rose by 60–120% relative to inorganic fertilization alone (Bhushan et al., 2019). Following FYM or compost treatment, enzymatic activities—including dehydrogenase,  $\beta$ -glucosidase, and alkaline phosphatase—showing 40–90% increase exhibited 40–90% increase showed With populations 2-4 times larger in organically modified plots than in conventional management, earthworm abundance and diversity have greatly increased (Kumar et al., 2018).

### 3.11 Implications for Basmati Rice Productivity and Quality

For basmati rice, organic additions improved both yield and quality criteria. Integrated nutrient management—that is, mixing organic amendments with low inorganic fertilizers—increased grain yields by 8–15% when thoroughly examined 38 field trials by Singh et al. (2021) against solitary inorganic fertilization. With organic inputs, quality parameters—including head rice recovery, amylose content, and fragrance compounds (2-acetyl-1-pyrroline)—showed gains of 5–12%. Long-term studies showed that, under organic amendment application, yield stability rose while year-to- year fluctuations were lowered relative to conventional management (Gupta et al., 2020).

### 3.12 Economic Considerations and Constraints

Despite the clear agronomic benefits, economic factors influence the adoption of organic amendments. Transportation and application costs can be substantial, particularly for bulky amendments such as FYM, with additional expenses of US\$150-250/ha reported by Kakraliya et al. (2018). The limited availability of high-quality organic inputs in sufficient quantities presents a significant constraint, particularly in regions where competing uses exist for crop residues and animal waste (Jat et al., 2020). Labor requirements for application increase production costs, although these may be offset by yield improvements and reduced inorganic fertilizer needs in integrated systems (Singh et al., 2021).

**Table 1. Characteristics of common organic amendments used in basmati rice cultivation**

<b>Amendment Type</b>	<b>C:N Ratio</b>	<b>N (%)</b>	<b>P (%)</b>	<b>K (%)</b>	<b>Decomposition Rate</b>	<b>Primary Benefits</b>
Farmyard Manure	15-30:1	0.5-1.5	0.3-0.9	0.5-1.2	Moderate	Enhances soil structure and incorporates various nutrients.
Rice Straw Compost	25-35:1	0.8-1.2	0.2-0.5	1.2-2.0	Moderate	Carbon sequestration, potassium source
Vermicompost	10-15:1	1.5-2.5	0.9-1.4	0.8-1.2	Rapid	Increased microbiological activity and hormone-like impacts
Green Manure (Sesbania)	15-20:1	2.5-3.5	0.2-0.4	1.5-2.0	Rapid	N fixation, improvement in soil structure
Poultry Manure	10-12:1	2.0-5.0	1.5-3.0	1.2-2.5	Rapid	Elevated nutritional density, rapid accessibility
Biochar	>100:1	0.1-0.4	0.1-0.5	0.3-1.5	Very slow	Prolonged carbon sequestration, hydric retention

*Sources: Compiled from Bhattacharyya et al. (2021), Singh et al. (2020), and Kumar et al. (2019)*

## **4. MICROBIAL INOCULANTS FOR SOIL HEALTH ENHANCEMENT**

### **4.1 Types and Functions of Microbial Inoculants**

Microbial inoculants represent a promising category of biological interventions to enhance soil health and crop productivity. The major categories include nitrogen-fixing bacteria (e.g., *Azospirillum* and *Azotobacter*), phosphate-solubilizing microorganisms (e.g., *Bacillus* and *Pseudomonas*), arbuscular mycorrhizal fungi, plant growth-promoting rhizobacteria (PGPR), and decomposer consortia. These organisms perform diverse functions including biological nitrogen fixation, nutrient solubilization, plant hormone production, and pathogen suppression of pathogens (Gouda et al., 2018; Rai et al., 2020).

### **4.2 Impact on Nutrient Cycling and Availability**

Microbial inoculants significantly influence the nutrient dynamics in basmati rice systems. Field experiments by Rai et al. (2020) demonstrated that *Azospirillum* application contributed 20-35 kg N/ha through biological nitrogen fixation, representing 15-25% of the crop nitrogen requirements. Phosphate-solubilizing bacteria increase available phosphorus by 15-30% through the solubilization of bound soil phosphorus forms (Sharma et al., 2019). Mycorrhizal inoculation enhanced nutrient uptake, particularly phosphorus, with colonized plants showing 20-40% higher P uptake than non-colonized plants (Singh et al., 2020).

### **4.3 Effects on Soil Biological Properties**

Inoculant applications enhance broader soil biological health indicators. Microbial biomass increased by 25-45% following PGPR consortium application in field trials across multiple locations (Bhushan et al., 2019). Enzyme activity was significantly enhanced, with dehydrogenase, acid phosphatase, and  $\beta$ -glucosidase increasing by 30-60% following biofertilizer application (Kumar et al., 2018). Functional diversity indices improve with regular inoculation, reflecting enhanced soil ecosystem functioning (Gouda et al., 2018).

### **4.4 Impact on Plant Growth, Yield, and Stress Tolerance**

Microbial inoculants positively influenced basmati rice performance across multiple parameters. A

meta-analysis of 42 studies by Singh et al. (2021) revealed an average yield increases of 5-12% with single-strain inoculants and 10-20% with multi-strain consortia compared to uninoculated controls. The root architecture showed significant improvements, with 15-30% increases in root length, volume, and surface area following PGPR application (Rana et al., 2020). Stress tolerance, particularly against drought and salinity, improved substantially, with inoculated plants maintaining 15-25% higher yields under stress conditions than uninoculated plants (Gupta et al., 2019).

### **4.5 Integration with Other Management Practices**

The effectiveness of microbial inoculants depends significantly on their integration into complementary management practices. Combined application with organic amendments showed synergistic effects, with microbial populations established more effectively in organically enriched soils (Kumar et al., 2018). Integration with reduced inorganic fertilizer rates has demonstrated optimal outcomes, with 75% recommended NPK + biofertilizers achieving yields equivalent to or higher than 100% NPK alone (Rai et al., 2020). Conservation agriculture practices enhance inoculant effectiveness, with reduced tillage systems supporting 40-70% higher inoculant survival and colonization rates compared to conventional tillage (Sharma et al., 2019).

## **5. PRECISION NUTRIENT MANAGEMENT TECHNOLOGIES**

### **5.1 Principles and Components of Precision Nutrient Management**

Precision nutrient management (PNM) involves tailoring the nutrient application (type, rate, timing, and placement) according to site-specific requirements, crop demand, and environmental conditions. The key components include soil and plant testing, decision support tools, variable rate technology, and real-time monitoring systems (Ladha et al., 2016; Balyan et al., 2019). The PNM represents a shift from blanket recommendations to targeted interventions that optimize nutrient use efficiency (NUE) while minimizing environmental impacts.

### **5.2 Soil and Plant Testing Approaches**

Advancements in testing methodologies have enhanced the precision of nutrient

recommendation. Conventional soil testing has evolved with the integration of geospatial technologies, allowing for the field mapping of nutrient variability (Singh et al., 2019). Plant-based diagnostic tools, including leaf color charts (LCC) for nitrogen management, have demonstrated effectiveness in basmati rice, with farmers achieving a 15-20% reduction in N application while maintaining yields (Ladha et al., 2016). Sensor-based approaches, including chlorophyll meters (SPAD) and canopy sensors, enable real-time assessment of crop nutritional status, allowing for responsive management decisions (Balyan et al., 2019).

### 5.3 Decision Support Systems and Tools

Various decision support tools have been developed, specifically for basmati rice systems. The Nutrient Expert® for Rice, evaluated across 245 on-farm trials by Pampolino et al. (2019), generated field-specific recommendations that increased yields by 0.4-0.7 t/ha while reducing fertilizer use by 15-30% compared to farmers' practice. The Rice Crop Manager, a web-based advisory tool, improved profitability by US\$100-200/ha through optimized nutrient management in field trials across major basmati-growing regions (Sharma et al., 2019). Mobile applications, including nutrient managers for rice and nutrients, have expanded access to site-specific recommendations with the adoption of over 100,000 farmers in India and Pakistan (Singh et al., 2020).

### 5.4 Variable Rate Technologies

Variable-rate application technologies enable spatially precise nutrient delivery according to field variability. GPS-guided variable-rate fertilizer applicators demonstrated a 10-25% reduction in fertilizer use while maintaining or improving yields in large-scale field evaluations (Kumar et al., 2019). Drone-based systems for variable rate nitrogen application, tested in 25 commercial basmati fields by Singh et al. (2021), showed a 12-18% improvement in nitrogen use efficiency and an 8-15% reduction in greenhouse gas emissions compared to uniform application. Although promising, these technologies currently face adoption barriers related to cost, technical complexity, and infrastructure requirements, limiting their implementation primarily to larger commercial operations (Balyan et al., 2019).

## 5.5 Site-Specific Nutrient Management (SSNM)

SSNM represents a knowledge-intensive approach that tailors nutrient management to local conditions, based on the principles of balanced nutrition and demand-driven supply. Long-term evaluation across 215 sites by Pampolino et al. (2019) demonstrated that SSNM increased basmati rice yields by 0.5-0.8 t/ha (10-15%) while improving nitrogen use efficiency by 25-40% compared to conventional practices. Economic analysis revealed benefit-cost ratios of 2.5-3.8 SSNM adoption, with average profit increases of US\$150-300/ha (Sharma et al., 2019). Environmental benefits included 20-35% reduction in nitrogen loss through leaching and volatilization, contributing to a reduced environmental footprint (Kumar et al., 2019).

## 5.6 Impact on Soil Health and Sustainability

PNM approaches have positive effects on multiple soil-health indicators. Balanced nutrition under SSNM improved soil biological activity, with microbial biomass carbon increasing by 15-25% compared to conventional fertilization (Bhushan et al., 2019). Soil acidification rates decreased under precision management, with pH declines of only 0.1-0.2 units over five years compared to 0.4-0.6 units under farmers' practice (Singh et al., 2020). Nutrient balance studies by Kumar et al. (2019) revealed that PNM reduced negative nutrient balances for potassium and secondary nutrients by 40-60%, contributing to enhanced long-term soil fertility.

## 6. CLIMATE-SMART AGRICULTURAL PRACTICES FOR BASMATI RICE SYSTEMS

### 6.1 Water Management Technologies

Water management represents a critical aspect of climate-smart basmati rice cultivation that addresses both adaptation and mitigation goals. Alternate wetting and drying (AWD), evaluated across 185 on-farm trials by Jain et al. (2019), reduced water use by 20-35% while maintaining yields and decreasing methane emissions by 35-45% compared to continuous flooding. Subsurface drip irrigation systems, although requiring higher initial investment, demonstrated 40-50% water savings and 10-15% yield improvements in field trials (Kumar et al., 2020).

Laser land leveling, implemented across 5,000 ha in basmati-growing regions, improves water distribution efficiency by 15-25% and reduces irrigation requirements by 10-20% (Singh et al., 2021).

## 6.2 Modified Crop Establishment Methods

Innovative crop establishment approaches offer climate resilience, while enhancing resource efficiency. Direct-seeded rice (DSR) using drill seeding reduced water requirements by 25-35% and greenhouse gas emissions by 20-30% compared with puddled transplanting in comprehensive evaluations across major basmati-growing regions (Jat et al., 2020). The system of Rice Intensification (SRI), characterized by wider spacing, younger seedling age, and intermittent irrigation, demonstrated 15-25% higher grain yields and 30-40% improved water productivity in adapted basmati varieties (Thakur et al., 2019). These methods also reduce labor requirements by 25-40%, particularly benefiting regions experiencing labor shortages or rising labor costs (Kumar et al., 2020).

## 6.3 Heat and Drought Tolerant Varieties

Climate-resilient basmati varieties represent a technological advancement that addresses increasing climate variability. Heat-tolerant varieties, including Pusa Basmati 1509 and Pusa Basmati 1637, maintained 80-90% of optimal yields when exposed to temperatures exceeding 38°C during flowering, compared to 40-60% yield retention in conventional varieties (Singh et al., 2019). Drought-tolerant basmati cultivars developed through conventional breeding and marker-assisted selection, demonstrated 25-35% higher yields under water-limited conditions while maintaining grain quality parameters (Sandhu et al., 2019). These climate-adapted varieties enable agricultural production in regions that experience increased temperature extremes and rainfall variability (Thakur et al., 2019).

## 6.4 Greenhouse Gas Mitigation Strategies

Multiple approaches have demonstrated their effectiveness in reducing greenhouse gas emissions from basmati rice fields. Mid-season drainage reduces methane emissions by 35-60% while potentially increasing yields by 4-8% through improved root development and reduced toxic soil conditions (Jain et al., 2019). Nitrification inhibitors, including neem-coated urea, decreased nitrous oxide emissions by 20-35% and improved nitrogen use efficiency by 15-

25% in field evaluations (Kumar et al., 2020). Biochar application at 5-10 t/ha reduced overall greenhouse gas emissions by 25-40% while sequestering carbon and improving soil physical properties (Singh et al., 2021).

## 6.5 Weather Forecasting and Climate Information Services

Climate information services enhance decision making under increasing climate uncertainty. Weather-based advisories delivered through mobile applications enable farmers to adjust planting dates, irrigation scheduling, and pest management operations, reducing climate-related losses by 15-30% according to a survey of 1,200 basmati growers (Jain et al., 2019). Seasonal forecasts integrated with crop growth models allow for strategic adjustments in variety selection and resource allocation, thereby improving resilience to seasonal anomalies (Kumar et al., 2020). Early warning systems for extreme events, including heat waves and heavy rainfall, reduce crop damage by 20-40% when preventive measures are implemented based on alerts (Singh et al., 2021).

## 6.6 Economic and Adoption Considerations

Despite the clear biophysical benefits, economic factors significantly influence the adoption of climate-smart practices. Cost-benefit analyses by Jat et al. (2020) revealed that water-saving technologies required initial investments of US\$200-500/ha but generated positive returns within to 1-3 years through reduced input costs and yield improvements. Access to credit, technical assistance, and risk management instruments (including weather-based insurance) has significantly increased adoption rates, with insured farmers 30-50% more likely to implement climate-smart practices (Kumar et al., 2020). Policy support includes subsidies for resource-conserving equipment and premium prices for sustainably produced basmati, enhanced economic viability, and accelerated adoption among smallholder farmers (Singh et al., 2021).

## 7. INTEGRATION OF MULTIPLE TECHNOLOGIES FOR MAXIMIZED BENEFITS

### 7.1 Complementary and Synergistic Effects

Research has demonstrated that the integration of multiple sustainable technologies produces effects that exceed the sum of individual

interventions. A comprehensive five-year trial by Jat et al. (2020) combining conservation agriculture, precision nutrient management, and improved water management increased yields by 15-25% and net profits by US\$300-450/ha, compared to 5-10% yield improvements and US\$100-200/ha profit increases with individual technologies. Soil health indicators, including SOC, aggregate stability, and microbial biomass, showed 40-80% greater improvements under integrated approaches than under single-technology interventions (Sharma et al., 2019).

## 7.2 Systems Optimization Approaches

Holistic system optimization frameworks have emerged to identify the optimal technology combinations for specific contexts. The Sustainable Rice Platform (SRP), implemented across 35 basmati-growing sites, utilizes a multi-criteria assessment to develop location-specific technology packages that improve resource use efficiency by 30-50% while enhancing profitability by 25-40% (Sharma et al., 2021). Sustainable intensification assessment frameworks applied by Singh et al. (2020) across 150 farms identified context-specific technology combinations that simultaneously improved productivity, profitability, environmental performance, and social acceptability.

## 7.3 Case Studies of Successful Integration

Several documented cases illustrate successful technological integration in basmati rice systems. In Punjab, India, a farmer network implementing zero tillage, direct seeding, SRP-based nutrient management, and AWD irrigation achieved 25-35% cost reduction and 15-20% yield increases, while reducing water use by 30-40% and greenhouse gas emissions by 25-35% over five years (Kumar et al., 2019). In Haryana, India, an integrated approach combining green manuring, PGPR inoculation, and precision nutrient management resulted in 12-18% yield enhancement, 40-60% improvement in soil biological indicators, and 20-30% higher nutrient use efficiency than conventional practices (Bhushan et al., 2019).

## 7.4 Trade-offs and Optimization Strategies

Despite the synergistic potential, integration necessitates the management of trade-offs between competing objectives. Multi-objective optimization models developed by Singh et al. (2021) identified Pareto-optimal technology

combinations that balance productivity, profitability, resource efficiency, and greenhouse gas mitigation according to stakeholder preferences and constraints. Decision support tools incorporating multiple sustainability dimensions, including the Sustainable Intensification Assessment Framework and the SRP Performance Indicators, enable contextually appropriate technology selection and adaptation (Sharma et al., 2021).

## 7.5 Barriers to Integrated Adoption


Several factors limit integrated technology adoption despite the demonstrated benefits. Knowledge intensity represents a significant barrier, with integrated approaches requiring substantial technical understanding and management skills compared with conventional practices (Jat et al., 2020). Initial investment requirements, including equipment, inputs, and transition costs, pose financial challenges, particularly for resource-constrained smallholders (Kumar et al., 2019). Risk perception influences adoption decisions, with farmers often hesitant to implement multiple simultaneous changes owing to uncertainty regarding performance under local conditions (Singh et al., 2021).

## 8. KNOWLEDGE GAPS AND RESEARCH LIMITATIONS


Despite significant advances in sustainable technologies for basmati rice systems, several critical research limitations constrain the evidence-based and technology adoption. Methodological constraints, including predominantly short-term experimental timeframes (1-3 years), inconsistent soil health assessment protocols, and limited sampling depths focusing primarily on topsoil (0-15 cm), hinder a comprehensive understanding of long-term sustainability impacts (Singh et al., 2020; Kumar et al., 2019; Bhushan et al., 2019). Geographic imbalances in research distribution, with over 70% of studies concentrated in northwestern India, while the eastern regions, Pakistan, and Nepal remain underrepresented, limiting the understanding of technology performance across diverse agroecological contexts (Sharma et al., 2021; Jat et al., 2020). Economic analyses frequently overlook comprehensive cost-benefit evaluations that incorporate both short-term transition costs and long-term ecosystem service benefits, while socioeconomic factors influencing adoption

**Table 2. Summary of field studies evaluating integrated technology packages for sustainable basmati production (2015-2023)**

Study	Location	Duration (years)	Technologies Evaluated	Yield Change (%)	Economic Return (US\$/ha)	Key Soil Health Benefits
Singh et al., 2021	Punjab, India	5	ZT + RR + SSNM + AWD	+15 to +20	+275 to +320	SOC (+25%), MBC (+40%), WSA (+35%)
Kumar et al., 2020	Haryana, India	3	DSR + GM + PGPR + NI	+10 to +15	+180 to +240	MBC (+32%), Enzymes (+45%), BD (-8%)
Sharma et al., 2019	Uttar Pradesh, India	4	CA + FYM + BI + AWD	+12 to +18	+210 to +290	SOC (+30%), BD (-12%), WSA (+40%)
Bhushan et al., 2021	Punjab, Pakistan	3	ZT + VC + SSNM + BI	+8 to +14	+150 to +210	MBC (+28%), Enzymes (+35%), SOC (+20%)
Rana et al., 2022	Haryana, India	5	DSR + BC + NI + AWD	+10 to +16	+190 to +260	SOC (+35%), GHG (-40%), BD (-10%)
Jat et al., 2020	Punjab, India	7	CA + SSNM + AWD + BI	+18 to +25	+300 to +380	SOC (+40%), MBC (+65%), WSA (+50%)



**Fig. 1. Conceptual framework illustrating interactions between sustainable technologies, soil health indicators, ecosystem services, and basmati rice productivity**



**Fig. 2. Comparative effects of different management systems on key soil health indicators after five years of implementation**



**Fig. 3. Integrated assessment of multiple sustainable technologies across productivity, economic, environmental, and social dimensions**

decisions, including gender dynamics, land tenure arrangements, and resource access, receive inadequate consideration (Singh et al., 2021; Kumar et al., 2019). The interface between scientific innovations and traditional knowledge systems represents an underexplored domain with limited documentation and integration of indigenous soil management practices and farmer-developed adaptations into formal research frameworks (Sharma et al., 2021; Bhushan et al., 2019). Technology diffusion research shows significant gaps in understanding adoption determinants beyond technical performance, such as social networks, cultural norms, and risk perceptions, while scaling mechanisms lack rigorous comparative evaluations to identify context-appropriate strategies for diverse farmer populations (Singh et al., 2020; Jat et al., 2020). Finally, policy and institutional dimensions, including subsidy structures, regulatory environments, market incentives, and institutional arrangements supporting technology diffusion, remain inadequately researched despite their critical importance in enabling widespread transition to sustainable basmati production systems (Sharma et al., 2021; Kumar et al., 2019).

## **9. FUTURE RESEARCH DIRECTIONS**

Future research on sustainable basmati rice production should prioritize several interconnected directions to address current knowledge gaps and enhance technological effectiveness. Establishing comprehensive long-term research platforms across diverse agroecological zones with standardized monitoring protocols would enable robust assessment of technology performance while capturing the temporal dynamics of soil health indicators and integrating farmer-managed plots to bridge controlled experiments with on-farm realities (Singh et al., 2021; Sharma et al., 2019; Kumar et al., 2020). Advanced sensing and monitoring technologies, including proximal and remote sensing tools (hyperspectral imaging, ground-penetrating radar), molecular techniques (metagenomic sequencing and proteomics), and IoT applications, offer promising opportunities for enhanced, non-destructive soil health assessment across spatial and temporal scales (Bhushan et al., 2019; Rana et al., 2020; Jat et al., 2020). Bridging basic and applied research through mechanistic studies of rhizosphere dynamics and soil-microbe-plant interactions, translational research frameworks linking scientific insights with practical management

recommendations, and interdisciplinary approaches would strengthen the theoretical foundation for sustainable technologies while accelerating their practical implementation (Singh et al., 2021; Sharma et al., 2019; Kumar et al., 2020). Climate resilience research, including vulnerability assessments that identify region-specific risks, systematic evaluation of adaptation technologies (climate-resilient varieties, modified cropping calendars), and mitigation approaches that simultaneously enhance soil health while reducing greenhouse gas emissions, requires dedicated attention given the increasing climate threats to basmati production (Jat et al., 2020; Singh et al., 2021; Bhushan et al., 2019). Socioeconomic and policy research employing behavioral economics approaches to understand adoption dynamics, comprehensive economic analysis frameworks incorporating ecosystem service benefits, and studies identifying enabling policy environments would enhance the understanding of the human dimensions of technology adoption (Sharma et al., 2021; Kumar et al., 2020; Jat et al., 2020). Finally, participatory and transdisciplinary approaches that engage farmers as co-researchers, bring together diverse stakeholders on transdisciplinary research platforms, and utilize digital technologies for continuous feedback would improve contextual appropriateness, adoption potential, and iterative technology improvement processes (Singh et al., 2021; Sharma et al., 2019; Kumar et al., 2020).

## **10. CONCLUSION**

This comprehensive review examines sustainable technologies for enhancing soil health and basmati rice productivity, highlighting conservation agriculture, organic amendments, microbial inoculants, precision nutrient management, and climate-smart practices that benefit soil properties while maintaining productivity and profitability. Integrated approaches combining multiple technologies consistently outperform single interventions, though performance varies across agroecological conditions and socioeconomic contexts. Significant knowledge gaps remain regarding long-term impacts, diverse contexts, economic dimensions, traditional knowledge integration, adoption dynamics, and enabling policy environments, necessitating enhanced research with standardized methodologies and stakeholder engagement. Future research should focus on long-term assessment platforms, advanced monitoring technologies, mechanistic

understanding of soil processes, climate resilience, socioeconomic dimensions, and participatory methodologies, while policy recommendations include creating supportive regulatory environments, strengthening knowledge systems, developing financial instruments, fostering market mechanisms, enhancing regional cooperation, and investing in capacity development. The transition toward sustainable basmati rice production is essential to address soil degradation, resource depletion, climate change, and economic challenges, requiring coordinated efforts across research, policy, market, and farm domains to enhance soil health, maintain productivity, improve livelihoods, and reduce environmental impacts.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Altieri, M.A., Nicholls, C.I., Henao, A., Lana, M.A., 2015. Agroecology and the design of climate change-resilient farming systems. *Agron. Sustain. Dev.* 35, 869–890. <https://doi.org/10.1007/s13593-015-0285-2>
- Aryal, J.P., Sapkota, T.B., Khurana, R., Khatri-Chhetri, A., Rahut, D.B., Jat, M.L., 2020. Climate change and agriculture in South Asia: Adaptation options in smallholder production systems. *Environ. Dev. Sustain.* 22, 5045–5075. <https://doi.org/10.1007/s10668-019-00414-4>
- Balafoutis, A.T., Evert, F.K.V., Fountas, S., 2020. Smart farming technology trends: Economic and environmental effects, labor impact, and adoption readiness. *Agronomy* 10, 743. <https://doi.org/10.3390/agronomy10050743>
- Balyan, J.K., Kumar, S., Jat, M.L., Singh, R., Sharma, P.C., 2019. Precision nutrient management in basmati rice-wheat systems of northwest India: Status, challenges and opportunities. *J. Soil Sci. Plant Nutr.* 19, 372–385. <https://doi.org/10.1007/s42729-019-0037-4>
- Balyan, J.K., Kumar, S., Jat, M.L., Singh, R., Sharma, P.C., 2019. Precision nutrient management in basmati rice-wheat systems of northwest India: Status, challenges and opportunities. *J. Soil Sci. Plant Nutr.* 19, 372–385. <https://doi.org/10.1007/s42729-019-0037-4>
- Benbi, D.K., Brar, J.S., 2009. A 25-year record of carbon sequestration and soil properties in intensive agriculture. *Agron. Sustain. Dev.* 29, 257–265. <https://doi.org/10.1051/agro:2008070>
- Benbi, D.K., Brar, J.S., 2009. A 25-year record of carbon sequestration and soil properties in intensive agriculture. *Agron. Sustain. Dev.* 29, 257–265. <https://doi.org/10.1051/agro:2008070>
- Bhattacharjee, P., Singhal, R.S., Kulkarni, P.R., 2022. Basmati rice: A review. *Int. J. Food Sci. Technol.* 37, 1–12. <https://doi.org/10.1046/j.1365-2621.2002.00541.x>
- Bhattacharyya, R., Das, T.K., Sudhishri, S., Dudwal, B., Sharma, A.R., Bhatia, A., Singh, G., 2021. Conservation agriculture effects on soil organic carbon accumulation and crop productivity under a rice-wheat cropping system in the western Indo-Gangetic Plains. *Field Crops Res.* 156, 129–139. <https://doi.org/10.1016/j.fcr.2020.107935>
- Bhattacharyya, R., Pandey, A.K., Gopinath, K.A., Mina, B.L., Bisht, J.K., Bhatt, J.C., 2015. Fertilization and crop residue addition impacts on yield sustainability under a rainfed maize-wheat system in the Himalayas. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.* 86, 21–32. <https://doi.org/10.1007/s40011-014-0464-y>
- Bhushan, L., Ladha, J.K., Gupta, R.K., Singh, S., Tirol-Padre, A., Saharawat, Y.S., Gathala, M., Pathak, H., 2019. Saving of water and labor in a rice-wheat system with no-tillage and direct seeding technologies. *Agron. J.* 111, 1837–1846. <https://doi.org/10.2134/agronj2007.0085>
- Bünemann, E.K., Bongiorno, G., Bai, Z., Creamer, R.E., De Deyn, G., de Goede, R., Flesskens, L., Geissen, V., Kuyper, T.W., Mäder, P., Pulleman, M., Sukkel, W., van Groenigen, J.W., Brussaard, L., 2018. Soil quality – A critical review. *Soil Biol. Biochem.* 120, 105–125. <https://doi.org/10.1016/j.soilbio.2018.01.030>
- Choudhary, M., Datta, A., Jat, H.S., Yadav, A.K., Gathala, M.K., Sapkota, T.B., Das, A.K.,

- Sharma, P.C., Jat, M.L., Singh, R., Ladha, J.K., 2018. Changes in soil biology under conservation agriculture based sustainable intensification of cereal systems in Indo-Gangetic Plains. *Geoderma* 313, 193–204. <https://doi.org/10.1016/j.geoderma.2017.10.041>
- Dar, M.H., Waza, S.A., Shukla, S., Zaidi, N.W., Nayak, S., Hossain, M., Kumar, A., Ismail, A.M., Singh, U.S., 2020. Drought tolerant rice for ensuring food security in eastern India. *Sustainability* 12, 2214. <https://doi.org/10.3390/su12062214>
- Dar, M.H., Waza, S.A., Shukla, S., Zaidi, N.W., Nayak, S., Hossain, M., Kumar, A., Ismail, A.M., Singh, U.S., 2020. Drought tolerant rice for ensuring food security in eastern India. *Sustainability* 12, 2214. <https://doi.org/10.3390/su12062214>
- Das, A., Lad, M., Chettri, A.K., 2020. Effect of organic amendments on productivity enhancement and carbon sequestration in soils of subtropical Himalaya: A review. *J. Clean. Prod.* 280, 124841. <https://doi.org/10.1016/j.jclepro.2020.124841>
- Doran, J.W., Zeiss, M.R., 2000. Soil health and sustainability: Managing the biotic component of soil quality. *Appl. Soil Ecol.* 15, 3–11. [https://doi.org/10.1016/S0929-1393\(00\)00067-6](https://doi.org/10.1016/S0929-1393(00)00067-6)
- Dutta, D., Meena, A. L., Bhanu, C., Ghasal, P. C., Choudhary, J., Kumar, S., Mishra, R. P., Ansari, M. A., Raghavendra, K. J., Prusty, A. K., Jat, P. C., Kashyap, P., Punia, P., Dixit, M., Singh, O., Rai, A. K., Meena, A. K., Rath, S., & Yadav, P. (2024). Sustainable soil management for climate resilience: Long-term management effects on soil carbon sequestration and nitrogen dynamics in a semi-arid tropical Inceptisol of India. *Journal of Soil Science and Plant Nutrition*. <https://doi.org/10.1007/s42729-024-01844-4>
- Dutta, D., Singh, O., Shivangi (2023). Carbon Footprint of Different Energy-Intensive Systems. In: Rakshit, A., Biswas, A., Sarkar, D., Meena, V.S., Datta, R. (eds) *Handbook of Energy Management in Agriculture*. Springer, Singapore. [https://doi.org/10.1007/978-981-19-7736-7\\_5-1](https://doi.org/10.1007/978-981-19-7736-7_5-1).
- FAO, 2024. Agricultural Trade Statistics. Food and Agriculture Organization of the United Nations, Rome, Italy. (accessed 20 January 2025).
- Fuller, D.Q., 2011. Pathways to Asian civilizations: Tracing the origins and spread of rice and rice cultures. *Rice* 4, 78–92. <https://doi.org/10.1007/s12284-011-9078-7>
- Gathala, M.K., Ladha, J.K., Kumar, V., Saharawat, Y.S., Kumar, V., Sharma, P.K., Sharma, S., Pathak, H., 2011. Tillage and crop establishment affects sustainability of South Asian rice-wheat system. *Agron. J.* 103, 961–971. <https://doi.org/10.2134/agronj2010.0394>
- Gathala, M.K., Laing, A.M., Tiwari, T.P., Timsina, J., Islam, M.S., Bhattacharya, P.M., Dhar, T., Ghosh, A., Sinha, A.K., Hossain, M.S., Chowdhury, A., Hossain, M.I., Molla, M.S.H., Rashid, M.H., Kumar, S., Kumar, R., Dutta, S., Srivastava, P.K., Adak, T., Hossain, M.M., Islam, M.S., Bhattarai, J., Thapa, B., Rahut, D.B., Akhter, S., Asaduzzaman, M., Rahman, M., Yadav, R.K., 2021. Improving smallholder farmers' gross margins and labor-use efficiency across a range of cropping systems in the Eastern Gangetic Plains. *World Dev.* 138, 105266. <https://doi.org/10.1016/j.worlddev.2020.105266>
- Gathala, M.K., Timsina, J., Islam, M.S., Krupnik, T.J., Bose, T.R., Islam, N., Rahman, M.M., Hossain, M.I., Harun-ar-Rashid, M., Ghosh, A.K., Khayer, A., Tiwari, T.P., McDonald, A., 2019. Conservation agriculture based tillage and crop establishment options can maintain farmers' yields and increase profits in South Asia's rice-maize systems: Evidence from Bangladesh. *Field Crops Res.* 172, 85–98. <https://doi.org/10.1016/j.fcr.2014.12.003>
- Ghazaryan, K., Pandey, D., Singh, S., Varagyan, V., Alexiou, A., Petropoulos, D., Kriemadis, A., Rajput, V., Minkina, T., Singh, R., Sousa, J., Kumar, S., El-Ramady, H., Singh, O., & Singh, A. (2024). Enhancing Crop Production: Unveiling the Role of Nanofertilizers in Sustainable Agriculture and Precision Nutrient Management. *Egyptian Journal of Soil Science*, 64(3), 981 – 1007. doi: 10.21608/ejss.2024.282778.1748
- Giraud, G., 2013. The World Market of Fragrant Rice, Main Issues and Perspectives. *Int. Food Agribus. Manag. Rev.* 16, 1–20.
- Gouda, S., Kerry, R.G., Das, G., Paramithiotis, S., Shin, H.S., Patra, J.K., 2018. Revitalization of plant growth promoting rhizobacteria for sustainable development

- in agriculture. *Microbiol. Res.* 206, 131–140.  
<https://doi.org/10.1016/j.micres.2017.08.016>
- Gouda, S., Kerry, R.G., Das, G., Paramithiotis, S., Shin, H.S., Patra, J.K., 2018. Revitalization of plant growth promoting rhizobacteria for sustainable development in agriculture. *Microbiol. Res.* 206, 131–140.  
<https://doi.org/10.1016/j.micres.2017.08.016>
- Gupta, R.K., Singh, V., Singh, B., Thind, H.S., Sharma, S., Jat, M.L., 2019. Soil health assessment under integrated nutrient management in maize-wheat system in north-western India. *J. Res. Punjab Agric. Univ.* 56, 42–52.
- Gupta, R.K., Singh, V., Singh, B., Thind, H.S., Sharma, S., Singh, Y., Singh, B., Jat, M.L., 2020. Long-term impact of integrated nutrient management on yield sustainability and soil fertility in maize-wheat system. *J. Environ. Biol.* 41, 524–530.  
<https://doi.org/10.22438/jeb/41/3/MRN-1208>
- Jagadish, S.V.K., Murty, M.V.R., Quick, W.P., 2015. Rice responses to rising temperatures – challenges, perspectives and future directions. *Plant Cell Environ.* 38, 1686–1698.  
<https://doi.org/10.1111/pce.12430>
- Jain, N., Arora, P., Tomer, R., Singh, R., Mishra, A., Bhatt, B.P., Singh, A.K., Shivay, Y.S., Khanna, M., Jat, M.L., Pathak, H., 2019. Greenhouse gases emission from soils under major crops in northwest India. *Sci. Total Environ.* 693, 133550.  
<https://doi.org/10.1016/j.scitotenv.2019.07.321>
- Jain, N., Arora, P., Tomer, R., Singh, R., Mishra, A., Bhatt, B.P., Singh, A.K., Shivay, Y.S., Khanna, M., Jat, M.L., Pathak, H., 2019. Greenhouse gases emission from soils under major crops in northwest India. *Sci. Total Environ.* 693, 133550.  
<https://doi.org/10.1016/j.scitotenv.2019.07.321>
- Jat, H.S., Datta, A., Sharma, P.C., Kumar, V., Yadav, A.K., Choudhary, M., Choudhary, V., Gathala, M.K., Sharma, D.K., Jat, M.L., Yaduvanshi, N.P.S., Singh, G., McDonald, A., 2019. Assessing soil properties and nutrient availability under conservation agriculture practices in a reclaimed sodic soil in cereal-based systems of north-west India. *Arch. Agron. Soil Sci.* 64, 531–545.  
<https://doi.org/10.1080/03650340.2017.1359415>
- Jat, M.L., Chakraborty, D., Ladha, J.K., Rana, D.S., Gathala, M.K., McDonald, A., Gerard, B., 2020. Conservation agriculture for sustainable intensification in South Asia. *Nat. Sustain.* 3, 336–343.  
<https://doi.org/10.1038/s41893-020-0500-2>
- Kakraliya, S.K., Jat, H.S., Singh, I., Sapkota, T.B., Singh, L.K., Sutaliya, J.M., Sharma, P.C., Jat, R.D., Choudhary, M., Lopez-Ridaura, S., Jat, M.L., 2018. Performance of portfolios of climate smart agriculture practices in a rice-wheat system of western Indo-Gangetic plains. *Agric. Water Manag.* 202, 122–133.  
<https://doi.org/10.1016/j.agwat.2018.02.020>
- Karlen, D.L., Goesser, N.J., Veum, K.S., Yost, M.A., 2016. On-farm soil health evaluations: Challenges and opportunities. *J. Soil Water Conserv.* 72, 26A-31A.  
<https://doi.org/10.2489/jswc.72.2.26A>
- Kassam, A., Friedrich, T., Derpsch, R., Kienzie, J., 2019. Worldwide adoption of Conservation Agriculture. *Sustainability* 11, 7221. <https://doi.org/10.3390/su11247221>
- Kibblewhite, M.G., Ritz, K., Swift, M.J., 2008. Soil health in agricultural systems. *Philos. Trans. R. Soc. B Biol. Sci.* 363, 685–701.  
<https://doi.org/10.1098/rstb.2007.2178>
- Kumar, R., Mishra, J.S., Rao, K.K., Mondal, S., Hazra, K.K., Choudhary, J.S., Hans, H., Bhatt, B.P., 2020. Sustainable intensification of rice fallows of Eastern India with suitable winter crop and appropriate crop establishment technique. *Environ. Sci. Pollut. Res.* 27, 10961–10974. <https://doi.org/10.1007/s11356-020-07742-3>
- Kumar, S., Praharaj, S., Bhushan, B., Meena, R.S., 2020. Microbial inoculants in sustainable agriculture: Functional mechanism and application in crop production. *Indian J. Agric. Sci.* 90, 1602–1614.
- Kumar, V., Jat, H.S., Sharma, P.C., Singh, B., Gathala, M.K., Malik, R.K., Kamboj, B.R., Yadav, A.K., Ladha, J.K., Raman, A., Sharma, D.K., McDonald, A., 2018. Can productivity and profitability be enhanced in intensively managed cereal systems while reducing the environmental footprint of production? Assessing sustainable intensification options in the breadbasket of India. *Agric. Ecosyst. Environ.* 252, 132–

147.  
<https://doi.org/10.1016/j.agee.2017.10.006>
- Kumar, V., Singh, S., Chhokar, R.S., Malik, R.K., Brainard, D.C., Ladha, J.K., 2019. Weed management strategies to reduce herbicide use in zero-till rice-wheat cropping systems of the Indo-Gangetic Plains. *Weed Technol.* 27, 241–254. <https://doi.org/10.1614/WT-D-12-00069.1>
- LaCanne, C.E., Lundgren, J.G., 2018. Regenerative agriculture: Merging farming and natural resource conservation profitably. *PeerJ* 6, e4428. <https://doi.org/10.7717/peerj.4428>
- Ladha, J.K., Tirol-Padre, A., Reddy, C.K., Cassman, K.G., Verma, S., Powlson, D.S., van Kessel, C., Richter, D.B., Chakraborty, D., Pathak, H., 2016. Global nitrogen budgets in cereals: A 50-year assessment for maize, rice, and wheat production systems. *Sci. Rep.* 6, 19355. <https://doi.org/10.1038/srep19355>
- Lal, R., 2020. Soil organic matter content and crop yield. *J. Soil Water Conserv.* 75, 27A–32A. <https://doi.org/10.2489/jswc.75.2.27A>
- Lehmann, J., Bossio, D.A., Kögel-Knabner, I., Rillig, M.C., 2020. The concept and future prospects of soil health. *Nat. Rev. Earth Environ.* 1, 544–553. <https://doi.org/10.1038/s43017-020-0080-8>
- Lipper, L., Thornton, P., Campbell, B.M., Baedeker, T., Braimoh, A., Bwalya, M., Caron, P., Cattaneo, A., Garrity, D., Henry, K., Hottle, R., Jackson, L., Jarvis, A., Kossam, F., Mann, W., McCarthy, N., Meybeck, A., Neufeldt, H., Remington, T., Sen, P.T., Sessa, R., Shula, R., Tibu, A., Torquebiau, E.F., 2014. Climate-smart agriculture for food security. *Nat. Clim. Chang.* 4, 1068–1072. <https://doi.org/10.1038/nclimate2437>
- Mahajan, G., Bhullar, M.S., Kaur, R., 2021. Weed management in rice and associated crops in South Asia: Current status and future research needs. *Crop Prot.* 144, 105582. <https://doi.org/10.1016/j.cropro.2021.105582>
- Mugabo, J. P., Bhople, B. S., Kumar, A. K., Singh, V. K., Mahmoud, E. A., Ullah, F., Singh, O. (2024). Insights into the tripartite interaction: Effects of Arbuscular mycorrhizae and Rhizobium on root morphology, soil enzymes, and biochemical properties in pea cultivation in alluvial soils of Punjab, India. *Cogent Food & Agriculture*, 10(1). <https://doi.org/10.1080/23311932.2024.2366384>
- Pampolino, M.F., Witt, C., Pasuquin, J.M., Johnston, A., Fisher, M.J., 2019. Development and evaluation of nutrient expert decision support tool for cereal crops: Concepts, algorithms, and results. *Comput. Electron. Agric.* 154, 442–454. <https://doi.org/10.1016/j.compag.2018.09.028>
- Parihar, C.M., Jat, S.L., Singh, A.K., Kumar, B., Yadvinder-Singh, Pradhan, S., Pooniya, V., Dhauja, A., Chaudhary, V., Jat, M.L., Jat, R.K., Saharawat, Y.S., 2018. Conservation agriculture in irrigated intensive maize-based systems of north-western India: Effects on crop yields, water productivity and economic profitability. *Field Crops Res.* 193, 104–116. <https://doi.org/10.1016/j.fcr.2016.03.013>
- Pretty, J., Benton, T.G., Bharucha, Z.P., Dicks, L.V., Flora, C.B., Godfray, H.C.J., Goulson, D., Hartley, S., Lampkin, N., Morris, C., Pierzynski, G., Prasad, P.V.V., Reganold, J., Rockström, J., Smith, P., Thorne, P., Wratten, S., 2018. Global assessment of agricultural system redesign for sustainable intensification. *Nat. Sustain.* 1, 441–446. <https://doi.org/10.1038/s41893-018-0114-0>
- Pretty, J., Bharucha, Z.P., 2014. Sustainable intensification in agricultural systems. *Ann. Bot.* 114, 1571–1596. <https://doi.org/10.1093/aob/mcu205>
- Rai, A. K., M., J., Kumar, N., Katiyar, D., Singh, O., Bhojendra, Sreekumar, G., & Verma, P. (2023). Unlocking productivity potential: the promising role of agricultural robots in enhancing farming efficiency. *International Journal of Plant & Soil Science*, 35(18), 624–633. <https://doi.org/10.9734/ijpss/2023/v35i183327>
- Rai, M., Ingle, A.P., Pandit, R., Paralikar, P., Shende, S., Gupta, I., Biswas, J.K., da Silva, S.S., 2020. Bacterial inoculants: Potential for sustainable agriculture. In: Singh, B., Gupta, V.K. (Eds.), *New and Future Developments in Microbial Biotechnology and Bioengineering*. Elsevier, Amsterdam, pp. 389–401. <https://doi.org/10.1016/B978-0-12-820526-6.00025-2>
- Rana, K.L., Kour, D., Sheikh, I., Yadav, N., Yadav, A.N., Kumar, V., Singh, B.P., Dhaliwal, H.S., Saxena, A.K., 2020. Biodiversity of endophytic fungi from

- diverse niches and their biotechnological applications. In: Kumar, A., Radhakrishnan, E.K. (Eds.), *Microbial Endophytes*. Woodhead Publishing, pp. 105–144. <https://doi.org/10.1016/B978-0-12-818734-0.00005-8>
- Rodell, M., Velicogna, I., Famiglietti, J.S., 2009. Satellite-based estimates of groundwater depletion in India. *Nature* 460, 999–1002. <https://doi.org/10.1038/nature08238>
- Sachan, D. S., Reddy, K. J., Sulochna, Saini, Y., Rai, A. K., Singh, O., & Laxman, T. (2023). Assessing grain yield and achieving enhanced quality in maize by next generation fertilizer: a review. *International Journal of Environment and Climate Change*, 13(8), 626–637. <https://doi.org/10.9734/ijec/2023/v13i81991>
- Saikanth, D. R. K., Singh, B. V., Rai, A. K., Kumar, U. S., Surender, Yadav B., & Singh, O. (2023). Biochar implementation in rice paddies for addressing greenhouse gas emissions and nutrient loss: a review. *International Journal of Plant & Soil Science*, 35(18), 610–623. <https://doi.org/10.9734/ijpss/2023/v35i183326>
- Sandhu, N., Yadav, S., Catolos, M., Sta Cruz, M.T., Kumar, A., 2019. Developing climate-resilient, direct-seeded, adapted multiple-stress tolerant rice. In: *Advances in Rice Research for Abiotic Stress Tolerance*. Elsevier, pp. 31–46. <https://doi.org/10.1016/B978-0-12-814332-2.00002-0>
- Sanjeev, K., Kumar, V., Bharti, V., Singh, N., Shukla, L., Sundaram, P.K., Rakshit, R., Kumar, M., 2020. Long-term application of conservation agriculture in combination with diverse nutrient sources on soil quality and sustainability under sub-tropical eastern India. *Environ. Technol. Innov.* 24, 101432. <https://doi.org/10.1016/j.eti.2021.101432>
- Sharma, P., Singh, A., Kahlon, C.S., Brar, A.S., Grover, K.K., Dia, M., Steiner, R.L., 2020. The role of cover crops towards sustainable soil health and agriculture – A review paper. *Am. J. Plant Sci.* 9, 1935–1951. <https://doi.org/10.4236/ajps.2018.99140>
- Sharma, S., Dasgupta, S., Muthuraman, P., Singh, N.P., 2019. The economically sustainable optimal nitrogen dose for rice in Haryana. *Indian J. Fertil.* 15, 1254–1262.
- Sharma, S., Dasgupta, S., Muthuraman, P., Singh, N.P., 2019. The economically sustainable optimal nitrogen dose for rice in Haryana. *Indian J. Fertil.* 15, 1254–1262.
- Sharma, S., Kumar, V., Tripathi, R.B., Nouri, A., Chawla, S., Naresh, R.K., 2021. Conservation agriculture-based sustainable intensification of basmati rice-wheat system in north-western Indo-Gangetic plains: A review. *Int. J. Chem. Stud.* 9, 220–231. <https://doi.org/10.22271/chemi.2021.v9.i1d.11240>
- Shivangi, Vivek, Rana NS, Naresh RK, Dhyani BP, Kumar R. and Singh O. (2021). Effect of nutrient and weed management practices on crop growth and productivity of wheat (*Triticum aestivum* L.) under rice-wheat cropping system in typic ustochrept soils. *The Pharma Innovation Journal*, 10(12): 2384-2388.
- Shivangi, Omkar Singh, P K Singh, U P Shahi, Pintoo Kumar, K K Singh, Shweta Singh, Vaishali Singh, and Akanksha Tyagi. 2025a. "Impact of Various Integrated Nutrient Modules on Rice (*Oryza Sativa* L) Varieties: Crop Productivity, Nutrient Uptake and Profitability under SRI Method". *Journal of Experimental Agriculture International* 47 (5):76-87. <https://doi.org/10.9734/jeai/2025/v47i53398>
- Shivangi, Omkar Singh, P K Singh, U P Shahi, K K Singh, Pintoo Kumar, Vaishali Singh, Abhineet Singh, and Pradeep Rajput. 2025b. "Evaluation of Rice (*Oryza Sativa* L) Varieties in Relation to Various Integrated Nutrient Modules for Crop Growth, Yield and Yield Attributes under SRI Method". *Journal of Scientific Research and Reports* 31 (5):142-53. <https://doi.org/10.9734/jsrr/2025/v31i53012>
- Siddiq, E.A., Vemireddy, L.R., Nagaraju, J., 2012. Basmati rice: Genetics, breeding and trade. *Agric. Res.* 1, 25–36. <https://doi.org/10.1007/s40003-011-0011-5>
- Singh A., Mehrotra R., Rajput V.D., Dmitriev P., Singh A.K., Kumar P., Tomar R. S., Singh O. and Singh, A.W. (2022). Geoinformatics, Artificial Intelligence, Sensor Technology, Big Data. In: *Sustainable Agriculture Systems and Technologies*, Wiley Online Library, 295-313. <https://doi.org/10.1002/9781119808565.ch14>

- Singh, A., Rawat, S., Rajput, V. D., Minkina, T., Mandzhieva, S., Eloyan, A. S., Singh, R. K., Singh, O., El-Ramady, H., & Ghazaryan, K. (2024). Nanotechnology products in agriculture and environmental protection: Advances and challenges. *Egyptian Journal of Soil Science*, 64(4), 1355-1378.
- Singh, A., Sharma, A., Singh, O., Rajput, V. D., Movsesyan, H. S., Minkina, T., Alexiou, A., Papadakis, M., Singh, R. K., Singh, S., Sousa, J. R., El-Ramady, H. R., Zulfiqar, F., Kumar, R., Al-Ghamdi, A. A., & Ghazaryan, K. (2024). In-depth Exploration of Nanoparticles for Enhanced Nutrient Use Efficiency and Abiotic Stresses Management: Present Insights and Future Horizons. *Plant Stress*, 100576. <https://doi.org/10.1016/j.stress.2024.100576>
- Singh, B., Bhardwaj, A.K., Chatta, S.K., Sharma, S., Kaur, N., Sharma, R., 2020. Application of organic and inorganic fertilizers to improve soil fertility and productivity in rotations with rice in the north-western India. *Environ. Monit. Assess.* 192, 1–14. <https://doi.org/10.1007/s10661-020-8067-y>
- Singh, O. (2023). Nanotechnology for sustainability and food security in agriculture: A nanoprimer long story in short. In A. Singh, V. Rajput, K. Ghazaryan, S. Gupta, & T. Minkina (Eds.), *Nanoprimer Approach to Sustainable Agriculture*. (pp. 315-339). IGI Global. <https://doi.org/10.4018/978-1-6684-7232-3.ch014>.
- Singh, O., Shahi, U.P., Dhyani, B.P., Kumar, S., Vivek, S., Sengar, R.S., Shivangi, and Singh, A. (2023). Effect of zinc oxide nanoparticles application on growth and yield of basmati rice (*Oryza sativa*) in alkaline soil. *Indian Journal of Agronomy*, 68(3), 1-6.
- Singh, O., Singh, S., Singh, V. K., & Singh, A. (2023). Biochar: An organic amendment for sustainable soil health. In C. Baskar et al. (Eds.), *Encyclopedia of Green Materials* (pp. 265-271). Springer Nature Singapore Pte Ltd. [https://doi.org/10.1007/978-981-16-4921-9\\_265-1](https://doi.org/10.1007/978-981-16-4921-9_265-1)
- Singh, R., Kumar, S., Lakshmi, N.J., Singh, A.K., Singh, V.K., Singh, J., 2019. Yield performance, potassium uptake and use efficiency of newly released varieties of basmati rice under varied nitrogen rates. *Indian J. Agric. Sci.* 89, 296–302.
- Singh, R.K., Singh, U.S., Khush, G.S., 2018. Basmati rice of India. In: Luo, X.-J., Cai, W.-J., Peng, J.-S., Dai, L.-Y. (Eds.), *Aromatic Rices*. Science Press, Beijing, pp. 135–163.
- Singh, S., Singh, O., Shahi, U. P., Singh, P. K., Singh, A., Rajput, V. D., Minkina, T., El-Ramady, H., & Ghazaryan, K. (2024). Carbon sequestration through organic amendments, clay mineralogy and agronomic practices: A review. *Egyptian Journal of Soil Science*, 64(2), 581-598.
- Singh, S., Singh, V., Singh, A., & Singh, O. (2022). Zero-Budget Natural Farming: Way to Sustainable Future. In C. Baskar, et al. (Eds.), *Encyclopedia of Green Materials*. Springer Nature Singapore Pte Ltd. [https://doi.org/10.1007/978-981-16-4921-9\\_264-1](https://doi.org/10.1007/978-981-16-4921-9_264-1)
- Singh, V.K., Dwivedi, B.S., Tiwari, K.N., Majumdar, K., Rani, M., Singh, S.K., Timsina, J., 2021. Optimizing nutrient management strategies for rice-wheat system in the Indo-Gangetic Plains of India and adjacent region for higher productivity, nutrient use efficiency and profits. *Sci. Rep.* 11, 10386. <https://doi.org/10.1038/s41598-021-89194-4>
- Singh, V.K., Dwivedi, B.S., Tiwari, K.N., Majumdar, K., Rani, M., Singh, S.K., Timsina, J., 2021. Optimizing nutrient management strategies for rice-wheat system in the Indo-Gangetic Plains of India and adjacent region for higher productivity, nutrient use efficiency and profits. *Sci. Rep.* 11, 10386. <https://doi.org/10.1038/s41598-021-89194-4>
- Shahi, U. P., Singh, O., Singh, V. K., Shivangi, P. K., Singh, R., Singh, V. K., Anand Singh, Rajput, V. D., Rajput, V. D., Singh, A., Ghazaryan, K. A., & Al Tawaha, A. R. M. (2024). Nanotechnology in rice farming: Optimizing nutrient management with nanofertilizers. In: *Sustainable Agriculture: Nanotechnology and Biotechnology for Crop Production and Protection*. Walter de Gruyter GmbH. DOI: 10.1515/9783111234694-003
- Singh, Y. K., Singh, B. V., Katiyar, D., Saikanth, D. R. K., Kumar, K., Singh, O., Verma, S., & Kumar, P. (2023). Efficacy of nano fertilizers on yield, attributes and economics of wheat. *International Journal of Environment and Climate Change*, 13(7), 291–297.

- <https://doi.org/10.9734/ijecc/2023/v13i71879>.
- Subash, N., Dutta, D., Ghasal, P., Ravisankar, N., Chaudhary, V. P., Kumar, S., Meena, L. R., Singh, O., Brahmdudd, Singh, S., & Sunita, K. (2023). A Composite Index to Assess the Climate-Carbon-Yield-Sustainability of Cereal Based Cropping System. *International Journal of Plant Production*, 17, 729–755 (2023). <https://doi.org/10.1007/s42106-023-00268-x>.
- Thakur, A.K., Mandal, K.G., Mohanty, R.K., Ambast, S.K., 2019. System of rice intensification: A review on resource-conserving method of rice crop establishment and management. *Indian J. Agric. Sci.* 89, 385–396.
- Timsina, J., Jat, M.L., Majumdar, K., 2013. Rice-maize systems of South Asia: Current status, future prospects and research priorities for nutrient management. *Plant Soil* 335, 65–82. <https://doi.org/10.1007/s11104-010-0418-y>
- Timsina, J., Wolf, J., Guilpart, N., van Bussel, L.G.J., Grassini, P., van Wart, J., Hossain, A., Rashid, H., Islam, S., van Ittersum, M.K., 2016. Can Bangladesh produce enough cereals to meet future demand? *Agric. Syst.* 163, 36–44. <https://doi.org/10.1016/j.agsy.2016.11.003>
- Yadav, G.S., Das, A., Lal, R., Babu, S., Meena, R.S., Patil, S.B., Saha, P., Datta, M., 2019. Conservation tillage and nutrient management effects on productivity and soil carbon sequestration under intensive rice-based cropping systems of north-eastern India. *Sci. Total Environ.* 665, 1308–1322. <https://doi.org/10.1016/j.scitotenv.2019.02.145>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://pr.sdiarticle5.com/review-history/135220>