



Archives of Current Research International

Volume 25, Issue 8, Page 412-436, 2025; Article no.ACRI.141230
ISSN: 2454-7077

A Study on Dual Hyperbolic Generalized Pandita Numbers

Fatih Zahid KALÇA^{a*} and Yüksel Soykan^a

^a *Department of Mathematics, Art and Science Faculty, Zonguldak Bülent Ecevit University, 67100, Zonguldak, Turkey.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/acri/2025/v25i81428>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/141230>

Original Research Article

Received: 28/05/2025

Published: 09/08/2025

ABSTRACT

This paper introduces the framework of generalized dual hyperbolic Pandita numbers, contributing a novel class of structured sequences to the expanding domain of number theory. Anchored in the principles of dual and hyperbolic systems, these constructs pave the way for exploring algebraic symmetries and recursive behaviors beyond classical formulations. Particular attention is devoted to notable special cases, including the dual hyperbolic Pandita and dual hyperbolic Pandita-Lucas numbers, whose properties are meticulously examined. To deepen understanding and facilitate computation, we derive explicit closed-form representations using Binet-type formulations, construct generating mechanisms through formal power series, and establish summative expressions with broad applicability. Additionally, matrix-based representations are developed to

**Corresponding author: E-mail: fzkakademi@hotmail.com;*

Cite as: KALÇA, Fatih Zahid, and Yüksel Soykan. 2025. "A Study on Dual Hyperbolic Generalized Pandita Numbers". *Archives of Current Research International* 25 (8):412-36. <https://doi.org/10.9734/acri/2025/v25i81428>.

offer an algebraic lens through which structural dynamics can be modeled and analyzed. These formulations not only enrich the theoretical foundations of discrete mathematics and symbolic computation but also highlight promising applications in engineering disciplines—particularly in the modeling of iterative systems, signal transformations, and the analysis of complex networks. The insights presented herein lay groundwork for future exploration into hybrid sequence systems and their role in interdisciplinary problem solving.

Keywords: Pandita numbers; Pandita-Lucas numbers; dual hyperbolic numbers; dual hyperbolic pandita numbers.

2020 Mathematics Subject Classification: 11B37, 11B39, 11B83.

1 INTRODUCTION

The hypercomplex numbers systems, (Kantor and Solodovnikov, 1989), are extensions of real numbers. Some commutative examples of hypercomplex number systems are complex numbers,

$$\mathbb{C} = \{z = a + ib : a, b \in \mathbb{R}, i^2 = -1\},$$

hyperbolic (double, split-complex) numbers, (Sobczyk, 1995),

$$\mathbb{H} = \{h = a + jb : a, b \in \mathbb{R}, j^2 = 1, j \neq \pm 1\},$$

and dual numbers, (Fjelstad and Gal, 1998),

$$\mathbb{D} = \{d = a + \varepsilon b : a, b \in \mathbb{R}, \varepsilon^2 = 0, \varepsilon \neq 0\}.$$

Some non-commutative examples of hypercomplex number systems are quaternions, (Hamilton, 1969),

$$\mathbb{H}_{\mathbb{Q}} = \{q = a_0 + ia_1 + ja_2 + ka_3 : a_0, a_1, a_2, a_3 \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\},$$

octonions (Baez, 2002) and sedenions (Soykan, 2019e) are part of a sequence of real algebras constructed through a recursive method known as the Cayley–Dickson process. The algebras \mathbb{C} (complex numbers), $\mathbb{H}_{\mathbb{Q}}$ (quaternions), \mathbb{O} (octonions) and \mathbb{S} (sedenions) are all derived from the real numbers \mathbb{R} via this doubling procedure. The process can be extended beyond sedenions to generate higher-dimensional algebras known as 2^n -ions [see for example (Biss et al., 2008; Imaeda and Imaeda, 2000; Moreno, 1998)].

Quaternions were introduced by the Irish mathematician W. R. Hamilton (1805–1865) as an extension of the complex numbers (Hamilton, 1969). Hyperbolic numbers with complex coefficients were first studied by J. Cockle in 1848 (Cockle, 1849). Later, H. H. Cheng and S. Thompson (Cheng and Thompson, 1996) introduced dual numbers with complex coefficients, which they termed complex dual numbers. Dual hyperbolic numbers were subsequently introduced by Akar et al. (2018).

A dual hyperbolic number is a hyper-complex number and is defined by

$$q = (a_0 + ja_1) + \varepsilon(a_2 + ja_3) = a_0 + ja_1 + \varepsilon a_2 + \varepsilon ja_3$$

where a_0, a_1, a_2 and a_3 are real numbers.

The set of all dual hyperbolic numbers are denoted by

$$\mathbb{H}_{\mathbb{D}} = \{a_0 + ja_1 + \varepsilon a_2 + \varepsilon ja_3 : a_0, a_1, a_2, a_3 \in \mathbb{R}, j^2 = 1, j \neq \pm 1, \varepsilon^2 = 0, \varepsilon \neq 0\}.$$

The base elements $\{1, j, \varepsilon, \varepsilon j\}$ of dual hyperbolic numbers satisfy the following properties (commutative multiplications):

$$\begin{aligned} 1.\varepsilon &= \varepsilon, 1.j = j, \varepsilon^2 = \varepsilon.\varepsilon = (\varepsilon\varepsilon)^2 = 0, j^2 = j.j = 1 \\ \varepsilon.j &= j.\varepsilon, \varepsilon.(\varepsilon j) = (\varepsilon j).\varepsilon = 0, j(\varepsilon j) = (\varepsilon j)j = \varepsilon \end{aligned}$$

where ε denotes the pure dual unit ($\varepsilon^2 = 0, \varepsilon \neq 0$), j denotes the hyperbolic unit ($j^2 = 1$), and εj denotes the dual hyperbolic unit ($(j\varepsilon)^2 = 0$).

The product of two dual hyperbolic numbers $q = a_0 + ja_1 + \varepsilon a_2 + j\varepsilon a_3$ and $p = b_0 + jb_1 + \varepsilon b_2 + j\varepsilon b_3$ is

$$qp = a_0b_0 + a_1b_1 + j(a_0b_1 + a_1b_0) + \varepsilon(a_0b_2 + a_2b_0 + a_1b_3 + a_3b_1) + j\varepsilon(a_0b_3 + a_1b_2 + a_2b_1 + b_0a_3)$$

and addition of dual hyperbolic numbers is defined as componentwise.

The set of dual hyperbolic numbers constitutes a commutative ring, a real vector space, and an algebra. However, H.D does not form a field, as not every dual hyperbolic number possesses a multiplicative inverse. For further details on the algebraic structure and properties of dual hyperbolic numbers, see (Akar et al., 2018).

We now recall the definition of generalized Pandita numbers.

A generalized Pandita sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, W_3)\}_{n \geq 0}$ is defined by the fourth-order recurrence relations

$$W_n = 2W_{n-1} - W_{n-2} + W_{n-3} - W_{n-4} \quad (1.1)$$

with the initial values W_0, W_1, W_2, W_3 not all being zero. The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = 2W_{-(n-1)} - W_{-(n-2)} + W_{-(n-3)} - W_{-(n-4)}$$

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1.1) holds for all integer n . Soykan has conducted a study on this particular sequence, for more details, see (Soykan, 2023).

The first few generalized Pandita numbers with positive subscript and negative subscript are given in the following Table 1.

Table 1. A few generalized Pandita numbers

n	W_n	W_{-n}
0	W_0	W_0
1	W_1	$W_0 - W_1 + 2W_2 - W_3$
2	W_2	$W_1 + W_2 - W_3$
3	W_3	$W_0 + W_1 - W_2$
4	$W_1 - W_0 - W_2 + 2W_3$	$2W_0 - 2W_1 + 2W_2 - W_3$
5	$W_1 - 2W_0 - W_2 + 3W_3$	$3W_2 - 2W_3$
6	$W_1 - 3W_0 - 2W_2 + 5W_3$	$3W_1 - 2W_2$
7	$2W_1 - 5W_0 - 4W_2 + 8W_3$	$3W_0 - 2W_1$
8	$3W_1 - 8W_0 - 6W_2 + 12W_3$	$W_0 - 3W_1 + 6W_2 - 3W_3$
9	$4W_1 - 12W_0 - 9W_2 + 18W_3$	$5W_1 - 2W_0 - W_2 - W_3$
10	$6W_1 - 18W_0 - 14W_2 + 27W_3$	$3W_0 + W_1 - 5W_2 + 2W_3$
11	$9W_1 - 27W_0 - 21W_2 + 40W_3$	$4W_0 - 8W_1 + 8W_2 - 3W_3$
12	$13W_1 - 40W_0 - 31W_2 + 59W_3$	$4W_1 - 4W_0 + 5W_2 - 4W_3$
13	$19W_1 - 59W_0 - 46W_2 + 87W_3$	$9W_1 - 12W_2 + 4W_3$

If we set $W_0 = 0, W_1 = 1, W_2 = 2, W_3 = 3$ then $\{W_n\}$ is the well-known Pandita sequence and if we set $W_0 = 4, W_1 = 2, W_2 = 2, W_3 = 5$ then $\{W_n\}$ is the well-known Pandita-Lucas sequence. In other words, Pandita sequence $\{P_n\}_{n \geq 0}$ and Pandita-Lucas sequence $\{S_n\}_{n \geq 0}$ are defined by the second-order recurrence relations

$$P_n = 2P_{n-1} - P_{n-2} + P_{n-3} - P_{n-4}, \quad P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 3, \quad n \geq 4, \quad (1.2)$$

and

$$S_n = 2S_{n-1} - S_{n-2} + S_{n-3} - S_{n-4}, \quad S_0 = 4, S_1 = 2, S_2 = 2, S_3 = 5, \quad n \geq 4. \quad (1.3)$$

The sequences $\{P_n\}_{n \geq 0}$ and $\{S_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$P_{-n} = P_{-(n-1)} - P_{-(n-2)} + 2P_{-(n-3)} - P_{-(n-4)}$$

and

$$S_{-n} = S_{-(n-1)} - S_{-(n-2)} + 2S_{-(n-3)} - S_{-(n-4)},$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrences (1.2) and (1.3) hold for all integer n .

We can list some important properties of generalized Pandita numbers that are needed.

- Binet formula of generalized Pandita sequence can be calculated using its characteristic equation which is given as

$$x^4 - 2x^3 + x^2 - x + 1 = (x^3 - x^2 - 1)(x - 1) = 0$$

The roots of characteristic equation are

$$\begin{aligned} \alpha &= \frac{1}{3} + \left(\frac{29}{54} + \sqrt{\frac{31}{108}} \right)^{1/3} + \left(\frac{29}{54} - \sqrt{\frac{31}{108}} \right)^{1/3}, \\ \beta &= \frac{1}{3} + \omega \left(\frac{29}{54} + \sqrt{\frac{31}{108}} \right)^{1/3} + \omega^2 \left(\frac{29}{54} - \sqrt{\frac{31}{108}} \right)^{1/3}, \\ \gamma &= \frac{1}{3} + \omega^2 \left(\frac{29}{54} + \sqrt{\frac{31}{108}} \right)^{1/3} + \omega \left(\frac{29}{54} - \sqrt{\frac{31}{108}} \right)^{1/3}, \\ \delta &= 1, \end{aligned}$$

where

$$\omega = \frac{-1 + i\sqrt{3}}{2} = \exp(2\pi i/3).$$

Using these roots and the recurrence relation, Binet formula can be given as

$$\begin{aligned} W_n &= \frac{z_1 \alpha^n}{3\alpha - 2} + \frac{z_2 \beta^n}{3\beta - 2} + \frac{z_3 \gamma^n}{3\gamma - 2} + z_4 \\ &= A_1 \alpha^n + A_2 \beta^n + A_3 \gamma^n + A_4, \end{aligned}$$

where z_1, z_2 and z_3 are given below

$$\begin{aligned} z_1 &= (\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0), \\ z_2 &= (\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0), \\ z_3 &= (\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0), \\ z_4 &= -W_3 + W_2 + W_0. \end{aligned}$$

and

$$\begin{aligned} A_1 &= \frac{z_1}{3\alpha - 2}, \\ A_2 &= \frac{z_2}{3\beta - 2}, \\ A_3 &= \frac{z_3}{3\gamma - 2}, \\ A_4 &= z_4. \end{aligned} \quad (1.4)$$

Binet formula of Pandita and Pandita-Lucas sequences are

$$P_n = \frac{\alpha^{n+3}}{3\alpha - 2} + \frac{\beta^{n+3}}{3\beta - 2} + \frac{\gamma^{n+3}}{3\gamma - 2} - 1,$$

and

$$S_n = \alpha^n + \beta^n + \gamma^n + 1,$$

respectively.

- The generating function for generalized Pandita numbers is

$$\sum_{n=0}^{\infty} W_n x^n = \frac{W_0 + (W_1 - 2W_0)x + (W_2 - 2W_1 + W_0)x^2 + (W_3 - 2W_2 + W_1 - W_0)x^3}{1 - 2x + x^2 - x^3 + x^4}.$$

For more details about generalized Pandita numbers, see (Soykan, 2023).

Next, we give the exponential generating function of $\sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ of the sequence W_n .

Lemma 1.1. (Kalça and Soykan, 2025), Lemma 1.4]. Suppose that $f_{W_n}(x) = \sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ is the exponential generating function of the generalized Pandita sequence $\{W_n\}$.

Then $\sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ is given by

$$\begin{aligned} \sum_{n=0}^{\infty} W_n \frac{x^n}{n!} &= \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)}{3\alpha - 2} e^{\alpha x} \\ &+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)}{3\beta - 2} e^{\beta x} \\ &+ \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)}{3\gamma - 2} e^{\gamma x} \\ &+ (-W_3 + W_2 + W_0)e^x. \end{aligned}$$

The previous Lemma 1.1 gives the following results as particular examples.

Corollary 1.2. Exponential generating function of Pandita and Pandita-Lucas numbers

$$\begin{aligned} \text{a)} \quad \sum_{n=0}^{\infty} P_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} \left(\frac{\alpha^{n+3}}{3\alpha - 2} + \frac{\beta^{n+3}}{3\beta - 2} + \frac{\gamma^{n+3}}{3\gamma - 2} - 1 \right) \frac{x^n}{n!} = \frac{\alpha^3 e^{\alpha x}}{3\alpha - 2} + \frac{\beta^3 e^{\beta x}}{3\beta - 2} + \frac{\gamma^3 e^{\gamma x}}{3\gamma - 2} - e^x. \\ \text{b)} \quad \sum_{n=0}^{\infty} S_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} (\alpha^n + \beta^n + \gamma^n + 1) \frac{x^n}{n!} = e^{\alpha x} + e^{\beta x} + e^{\gamma x} + e^x. \end{aligned}$$

Next, we give some information on published papers related to hyperbolic and dual hyperbolic numbers in literature.

- Cockle (1849) presented the hyperbolic numbers with complex coefficients.
- Akar et al. (2018) introduced the dual hyperbolic numbers.
- Cheng and Thompson (1996) studied dual numbers with complex coefficients.

Next, we give some information related to dual hyperbolic sequences presented in literature.

- Soykan et al. (2021) introduced dual hyperbolic generalized Pell numbers given by

$$\widehat{V}_n = V_n + jV_{n+1} + \varepsilon V_{n+2} + j\varepsilon V_{n+3}$$

where generalized Pell numbers are given by $V_n = 2V_{n-1} + V_{n-2}$, $V_0 = a$, $V_1 = b$ ($n \geq 2$) with the initial values V_0, V_1 not all being zero.

- Cihan et al. (2019) studied dual hyperbolic Fibonacci and Lucas numbers given by, respectively,

$$DHF_n = F_n + jF_{n+1} + \varepsilon F_{n+2} + j\varepsilon F_{n+3},$$

$$DHL_n = L_n + jL_{n+1} + \varepsilon L_{n+2} + j\varepsilon L_{n+3}.$$

where Fibonacci and Lucas numbers, respectively, given by $F_n = F_{n-1} + F_{n-2}$, $F_0 = 0$, $F_1 = 1$, $L_n = L_{n-1} + L_{n-2}$, $L_0 = 2$, $L_1 = 1$.

- Soykan et al. (2023) introduced dual hyperbolic generalized Jacopsthal numbers given by

$$\widehat{J}_n = J_n + jJ_{n+1} + \varepsilon J_{n+2} + j\varepsilon J_{n+3}$$

where $J_n = J_{n-1} + 2J_{n-2}$, $J_0 = a$, $J_1 = b$.

- Bród et al. (2020) studied dual hyperbolic generalized Balancing numbers are

$$DHB_n = B_n + jB_{n+1} + \varepsilon B_{n+2} + j\varepsilon B_{n+3}$$

where $B_n = 6B_{n-1} - B_{n-2}$, $B_0 = 0$, $B_1 = 1$.

- Yılmaz and Soykan (2024) introduced dual hyperbolic generalized Guglielmo numbers are

$$\widehat{T}_0 = T_0 + jT_1 + \varepsilon T_2 + j\varepsilon T_3$$

where $T_n = 3T_{n-1} - 3T_{n-2} + T_{n-3}$, $T_0 = 0$, $T_1 = 1$, $T_2 = 3$.

- Dikmen (2025) introduced dual hyperbolic generalised Leonardo numbers given by

$$\widehat{l}_0 = l_0 + jl_1 + \varepsilon l_2 + j\varepsilon l_3$$

$l_n = 2l_{n-1} - l_{n-3}$, $l_0 = 1$, $l_1 = 1$, $l_2 = 3$.

- Soykan et al. (2021) introduced dual hyperbolic generalized Woodall numbers given by

$$\widehat{R}_0 = R_0 + jR_1 + \varepsilon R_2 + j\varepsilon R_3$$

where $R_n = 5R_{n-1} - 8R_{n-2} + 4R_{n-3}$, $R_0 = -1$, $R_1 = 1$, $R_2 = 7$.

In this paper, we define the dual hyperbolic generalized Pandita numbers in the next section and give some properties of them.

2 DUAL HYPERBOLIC GENERALIZED PANDITA NUMBERS AND THEIR GENERATING FUNCTIONS AND BINET'S FORMULAS

In this section, we define dual hyperbolic generalized Pandita numbers and present generating functions and Binet formulas for them. We now define dual hyperbolic generalized Pandita numbers over \mathbb{H}_D . The n th dual hyperbolic generalized Pandita number is

$$\widehat{W}_n = W_n + jW_{n+1} + \varepsilon W_{n+2} + j\varepsilon W_{n+3}. \quad (2.1)$$

The sequence $\{\widehat{W}_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$\widehat{W}_{-n} = W_{-n} + jW_{-n+1} + \varepsilon W_{-n+2} + j\varepsilon W_{-n+3}.$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrence (2.2) holds for all integer n . Note that

$$\begin{aligned} \widehat{W}_0 &= W_0 + jW_1 + \varepsilon W_2 + j\varepsilon W_3 \\ \widehat{W}_1 &= W_1 + jW_2 + \varepsilon W_3 + j\varepsilon W_4 = W_1 + jW_2 + \varepsilon W_3 + j\varepsilon(W_1 - W_0 - W_2 + 2W_3) \\ \widehat{W}_2 &= W_2 + jW_3 + \varepsilon W_4 + j\varepsilon W_5 = W_2 + jW_3 + \varepsilon(W_1 - W_0 - W_2 + 2W_3) + j\varepsilon(W_1 - 2W_0 - W_2 + 3W_3) \\ \widehat{W}_3 &= W_3 + jW_4 + \varepsilon W_5 + j\varepsilon W_6 = W_3 + j(W_1 - W_0 - W_2 + 2W_3) + \varepsilon(W_1 - 2W_0 - W_2 + 3W_3) \\ &\quad + j\varepsilon(W_1 - 3W_0 - 2W_2 + 5W_3) \end{aligned}$$

It can be easily shown that

$$\widehat{W}_n = 2\widehat{W}_{n-1} - \widehat{W}_{n-2} + \widehat{W}_{n-3} - \widehat{W}_{n-4} \quad (2.2)$$

and

$$\widehat{W}_{-n} = \widehat{W}_{-(n-1)} - \widehat{W}_{-(n-2)} + 2\widehat{W}_{-(n-3)} - \widehat{W}_{-(n-4)}$$

The first few dual hyperbolic generalized Pandita numbers with positive subscript and negative subscript are given in the following Table 2.

Table 2. A few dual hyperbolic generalized Pandita numbers

n	\widehat{W}_n	\widehat{W}_{-n}
0	\widehat{W}_0	\widehat{W}_0
1	\widehat{W}_1	$\widehat{W}_0 - \widehat{W}_1 + 2\widehat{W}_2 - \widehat{W}_3$
2	\widehat{W}_2	$\widehat{W}_1 + \widehat{W}_2 - \widehat{W}_3$
3	\widehat{W}_3	$\widehat{W}_0 + \widehat{W}_1 - \widehat{W}_2$
4	$\widehat{W}_1 - \widehat{W}_0 - \widehat{W}_2 + 2\widehat{W}_3$	$2\widehat{W}_0 - 2\widehat{W}_1 + 2\widehat{W}_2 - \widehat{W}_3$
5	$\widehat{W}_1 - 2\widehat{W}_0 - \widehat{W}_2 + 3\widehat{W}_3$	$3\widehat{W}_2 - 2\widehat{W}_3$
6	$\widehat{W}_1 - 3\widehat{W}_0 - 2\widehat{W}_2 + 5\widehat{W}_3$	$3\widehat{W}_1 - 2\widehat{W}_2$
7	$2\widehat{W}_1 - 5\widehat{W}_0 - 4\widehat{W}_2 + 8\widehat{W}_3$	$3\widehat{W}_0 - 2\widehat{W}_1$
8	$3\widehat{W}_1 - 8\widehat{W}_0 - 6\widehat{W}_2 + 12\widehat{W}_3$	$\widehat{W}_0 - 3\widehat{W}_1 + 6\widehat{W}_2 - 3\widehat{W}_3$
9	$4\widehat{W}_1 - 12\widehat{W}_0 - 9\widehat{W}_2 + 18\widehat{W}_3$	$5\widehat{W}_1 - 2\widehat{W}_0 - \widehat{W}_2 - \widehat{W}_3$
10	$6\widehat{W}_1 - 18\widehat{W}_0 - 14\widehat{W}_2 + 27\widehat{W}_3$	$3\widehat{W}_0 + \widehat{W}_1 - 5\widehat{W}_2 + 2\widehat{W}_3$
11	$9\widehat{W}_1 - 27\widehat{W}_0 - 21\widehat{W}_2 + 40\widehat{W}_3$	$4\widehat{W}_0 - 8\widehat{W}_1 + 8\widehat{W}_2 - 3\widehat{W}_3$
12	$13\widehat{W}_1 - 40\widehat{W}_0 - 31\widehat{W}_2 + 59\widehat{W}_3$	$4\widehat{W}_1 - 4\widehat{W}_0 + 5\widehat{W}_2 - 4\widehat{W}_3$
13	$19\widehat{W}_1 - 59\widehat{W}_0 - 46\widehat{W}_2 + 87\widehat{W}_3$	$9\widehat{W}_1 - 12\widehat{W}_2 + 4\widehat{W}_3$

As special cases, the n th dual hyperbolic Pandita numbers and the n th dual hyperbolic Pandita-Lucas numbers are given as

$$\widehat{P}_n = P_n + jP_{n+1} + \varepsilon P_{n+2} + j\varepsilon P_{n+3} \quad (2.3)$$

and

$$\widehat{S}_n = S_n + jS_{n+1} + \varepsilon S_{n+2} + j\varepsilon S_{n+3} \quad (2.4)$$

respectively. The sequences $\{\widehat{P}_n\}_{n \geq 0}$ and $\{\widehat{S}_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$\widehat{P}_{-n} = P_{-(n-1)} - P_{-(n-2)} + 2P_{-(n-3)} - P_{-(n-4)}$$

and

$$\widehat{S}_{-n} = S_{-(n-1)} - S_{-(n-2)} + 2S_{-(n-3)} - S_{-(n-4)}$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrence (2.3) and (2.4) holds for all integer n

For dual hyperbolic Pandita numbers (taking $W_n = P_n$, $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 3$), we get

$$\begin{aligned} \widehat{P}_0 &= j + 2\varepsilon + 3j\varepsilon, \\ \widehat{P}_1 &= 2j + 3\varepsilon + 5j\varepsilon + 1, \\ \widehat{P}_2 &= 3j + 5\varepsilon + 8j\varepsilon + 2, \end{aligned}$$

and for dual hyperbolic Pandita-Lucas numbers (taking $W_n = S_n$, $S_0 = 4, S_1 = 2, S_2 = 2, S_3 = 5$), we get

$$\begin{aligned} \widehat{S}_0 &= 2j + 2\varepsilon + 5j\varepsilon + 4, \\ \widehat{S}_1 &= 2j + 5\varepsilon + 6j\varepsilon + 2, \\ \widehat{S}_2 &= 5j + 6\varepsilon + 7j\varepsilon + 2 \end{aligned}$$

A few dual hyperbolic Pandita numbers and dual hyperbolic Pandita-Lucas numbers with positive subscript and negative subscript are given in the following Table 3 and Table 4.

Table 3. Dual hyperbolic Pandita numbers

n	\widehat{P}_n	\widehat{P}_{-n}
0	$j + 2\varepsilon + 3j\varepsilon$	$j + 2\varepsilon + 3j\varepsilon$
1	$2j + 3\varepsilon + 5j\varepsilon + 1$	$\varepsilon + 2j\varepsilon$
2	$3j + 5\varepsilon + 8j\varepsilon + 2$	$-j\varepsilon$
3	$5j + 8\varepsilon + 12j\varepsilon + 3$	-1
4	$8j + 12\varepsilon + 18j\varepsilon + 5$	$-j - 1$
5	$12j + 18\varepsilon + 27j\varepsilon + 8$	$-j - \varepsilon$

Table 4. Dual hyperbolic Pandita-Lucas numbers

n	\widehat{S}_n	\widehat{S}_{-n}
0	$2j + 2\varepsilon + 5j\varepsilon + 4$	$2j + 2\varepsilon + 5j\varepsilon + 4$
1	$2j + 5\varepsilon + 6j\varepsilon + 2$	$-4j + 2\varepsilon + 2j\varepsilon + 1$
2	$5j + 6\varepsilon + 7j\varepsilon + 2$	$j + 4\varepsilon + 2j\varepsilon - 1$
3	$6j + 7\varepsilon + 11j\varepsilon + 5$	$\varepsilon - j + 4j\varepsilon + 4$
4	$7j + 11\varepsilon + 16j\varepsilon + 6$	$4j - \varepsilon + j\varepsilon + 3$
5	$11j + 16\varepsilon + 22j\varepsilon + 7$	$-3j + 4\varepsilon - j\varepsilon - 4$

Now, we will state Binet's formula for the dual hyperbolic generalized Pandita numbers and in the rest of the paper, we fix the following notations:

$$\hat{\alpha} = 1 + j\alpha + \varepsilon\alpha^2 + j\varepsilon\alpha^3, \quad (2.5)$$

$$\hat{\beta} = 1 + j\beta + \varepsilon\beta^2 + j\varepsilon\beta^3. \quad (2.6)$$

$$\hat{\gamma} = 1 + j\gamma + \varepsilon\gamma^2 + j\varepsilon\gamma^3 \quad (2.7)$$

$$\hat{\delta} = \hat{1} = 1 + j + \varepsilon + j\varepsilon, \quad (2.8)$$

Note that we have the following identities:

$$\begin{aligned} \hat{\alpha}^2 &= 1 + \alpha^2 + 2\alpha j + 2\alpha^2 (\alpha^2 + 1) \varepsilon + 4\alpha^3 j\varepsilon \\ \hat{\beta}^2 &= 1 + \beta^2 + 2j\beta + (2\beta^4 + 2\beta^2)\varepsilon + 4j\varepsilon\beta^3, \\ \hat{\alpha}\hat{\beta} &= 1 + \alpha\beta + (\alpha + \beta)j + (\alpha^2 + \beta^2 + 2\alpha\beta^3 + \alpha^3\beta)\varepsilon + (\alpha + \beta)(\alpha^2 + \beta^2)j\varepsilon, \\ \hat{\alpha}^2\hat{\beta} &= 1 + \alpha^2 + \beta^2 + \alpha^2\beta^2 + 2(\alpha\beta + 1)(\alpha + \beta)j + 2(\alpha^2 + \beta^2 + \alpha^2\beta^2 + 4\alpha\beta + 1)(\alpha^2 + \beta^2)\varepsilon \\ &\quad + 4(\alpha + \beta)(\alpha^2 + \beta^2 + \alpha\beta^3)j\varepsilon, \\ \hat{\alpha}\hat{\beta}^2 &= 1 + \beta^2 + 2\alpha\beta + (\alpha + 2\beta + \alpha\beta^2)j + (\beta^2 + 2\alpha\beta + 1)(\alpha^2 + 2\beta^2)\varepsilon + (\alpha + 2\beta + \alpha\beta^2)(\alpha^2 + 2\beta^2)j\varepsilon, \\ \hat{\alpha}^2\hat{\beta}^2 &= 1 + \beta^2 + \alpha^2 + \alpha^2\beta^2 + 4\alpha\beta + 2(\alpha\beta + 1)(\alpha + \beta)j + 2(\alpha^2 + \beta^2 + \alpha^2\beta^2 + 4\alpha\beta + 1)(\alpha^2 + \beta^2)\varepsilon \\ &\quad + 4(\alpha\beta + 1)(\alpha + \beta)(\alpha^2 + \beta^2)j\varepsilon \end{aligned}$$

Theorem 2.1. (Binet's Formula) For any integer n , the n th dual hyperbolic generalized Pandita number is

$$\widehat{W}_n = A_1\alpha^n\hat{\alpha} + A_2\beta^n\hat{\beta} + A_3\gamma^n\hat{\gamma} + \hat{1}A_4. \quad (2.9)$$

where $\hat{\alpha}, \hat{\beta}, \hat{\gamma}, \hat{\delta}$ are given as (2.5)-(2.8)

Proof. Using Binet's formula

$$W_n = A_1\alpha^n + A_2\beta^n + A_3\gamma^n + A_4.$$

where A_1, A_2, A_3, A_4 are given in (1.4) we get

$$\begin{aligned} \widehat{W}_n &= W_n + jW_{n+1} + \varepsilon W_{n+2} + j\varepsilon W_{n+3} \\ &= A_1\alpha^n + A_2\beta^n + A_3\gamma^n + A_4 + j(A_1\alpha^{n+1} + A_2\beta^{n+1} + A_3\gamma^{n+1} + A_4) \\ &\quad + \varepsilon(A_1\alpha^{n+2} + A_2\beta^{n+2} + A_3\gamma^{n+2} + A_4) + j\varepsilon(A_1\alpha^{n+3} + A_2\beta^{n+3} + A_3\gamma^{n+3} + A_4) \\ &= A_1\alpha^n(1 + j\alpha + \varepsilon\alpha^2 + j\varepsilon\alpha^3) + A_2\beta^n(1 + j\beta + \varepsilon\beta^2 + j\varepsilon\beta^3) \\ &\quad + A_3\gamma^n(1 + j\gamma + \varepsilon\gamma^2 + j\varepsilon\gamma^3) + A_4(1 + j + \varepsilon + j\varepsilon) \\ &= A_1\alpha^n\hat{\alpha} + A_2\beta^n\hat{\beta} + A_3\gamma^n\hat{\gamma} + \hat{1}A_4. \end{aligned}$$

This proves (2.9). \square

As special cases, for any integer n , the Binet's Formula of n th dual hyperbolic Pandita number is

$$\hat{P}_n = \frac{\alpha^{n+3}\hat{\alpha}}{3\alpha - 2} + \frac{\beta^{n+3}\hat{\beta}}{3\beta - 2} + \frac{\gamma^{n+3}\hat{\gamma}}{3\gamma - 2} - \hat{1} \quad (2.10)$$

and the Binet's Formula of n th dual hyperbolic Pandita-Lucas number is

$$\hat{S}_n = \hat{\alpha}\alpha^n + \hat{\beta}\beta^n + \hat{\gamma}\gamma^n + \hat{1}, \quad (2.11)$$

Next, we present generating function.

Theorem 2.2. Let $f_{\widehat{W}_n}(x) = \sum_{n=0}^{\infty} \widehat{W}_n x^n$ denote the generating function of dual hyperbolic generalized Pandita numbers is given as follows:

$$f_{\widehat{W}_n}(x) = \sum_{n=0}^{\infty} \widehat{W}_n x^n = \frac{\widehat{W}_0 + (\widehat{W}_1 - 2\widehat{W}_0)x + (\widehat{W}_2 - 2\widehat{W}_1 + \widehat{W}_0)x^2 + (\widehat{W}_3 - 2\widehat{W}_2 + \widehat{W}_1 - \widehat{W}_0)x^3}{1 - 2x + x^2 - x^3 + x^4}.$$

Proof. Using the definition of dual hyperbolic Pandita numbers, and subtracting $xf(x)$, $x^2f(x)$ and $x^3f(x)$ from $f(x)$ we obtain $(1 - 2x + x^2 - x^3 + x^4)f_{\widehat{W}_n}(x)$

$$\begin{aligned} & (1 - 2x + x^2 - x^3 + x^4)f_{\widehat{W}_n}(x) \\ &= \sum_{n=0}^{\infty} \widehat{W}_n x^n - 2x \sum_{n=0}^{\infty} \widehat{W}_n x^n + x^2 \sum_{n=0}^{\infty} \widehat{W}_n x^n - x^3 \sum_{n=0}^{\infty} \widehat{W}_n x^n + x^4 \sum_{n=0}^{\infty} \widehat{W}_n x^n, \\ &= \sum_{n=0}^{\infty} \widehat{W}_n x^n - 2 \sum_{n=0}^{\infty} \widehat{W}_n x^{n+1} + \sum_{n=0}^{\infty} \widehat{W}_n x^{n+2} - \sum_{n=0}^{\infty} \widehat{W}_n x^{n+3} + \sum_{n=0}^{\infty} \widehat{W}_n x^{n+4}, \\ &= \sum_{n=0}^{\infty} \widehat{W}_n x^n - 2 \sum_{n=1}^{\infty} \widehat{W}_{(n-1)} x^n + \sum_{n=2}^{\infty} \widehat{W}_{(n-2)} x^n - \sum_{n=3}^{\infty} \widehat{W}_{(n-3)} x^n + \sum_{n=4}^{\infty} \widehat{W}_{(n-4)} x^n, \\ &= (\widehat{W}_0 + \widehat{W}_1 x + \widehat{W}_2 x^2 + \widehat{W}_3 x^3) - 2(\widehat{W}_0 x + \widehat{W}_1 x^2 + \widehat{W}_2 x^3) + (\widehat{W}_0 x^2 + \widehat{W}_1 x^3) - \widehat{W}_0 x^3 \\ &\quad + \sum_{n=4}^{\infty} (\widehat{W}_n - 2\widehat{W}_{n-1} - \widehat{W}_{n-2} - \widehat{W}_{n-3} + \widehat{W}_{n-4}) x^n, \\ &= \widehat{W}_0 + (\widehat{W}_1 - 2\widehat{W}_0)x + (\widehat{W}_2 - 2\widehat{W}_1 + \widehat{W}_0)x^2 + (\widehat{W}_3 - 2\widehat{W}_2 + \widehat{W}_1 - \widehat{W}_0)x^3. \end{aligned}$$

And rearranging above equation, we get (2.2). \square

The following results are immediate consequences of the preceding Theorem.

Corollary 2.3. For all integers n , we have following identities:

$$\begin{aligned} \text{a) } \sum_{n=0}^{\infty} \widehat{P}_n x^n &= \frac{(j + 5\varepsilon + 4j\varepsilon) + (1 - \varepsilon - j\varepsilon)x + (\varepsilon + j\varepsilon)x^2 + (3j\varepsilon)x^3}{1 - 2x + x^2 - x^3 + x^4}, \\ \text{b) } \sum_{n=0}^{\infty} \widehat{S}_n x^n &= \frac{(2j + 2\varepsilon + 5j\varepsilon + 4) + (\varepsilon - 2j - 6x - 4j\varepsilon)x + (3j - 2\varepsilon + 2)x^2 + (2\varepsilon - 4j + 8j\varepsilon + 7)x^3}{1 - 2x + x^2 - x^3 + x^4}. \end{aligned}$$

Theorem (2.2) gives the following results as special cases,

$$\begin{aligned} (1 - 2x + x^2 - x^3 + x^4)f_{\widehat{P}_n}(x) &= \widehat{P}_0 + (\widehat{P}_1 - 2\widehat{P}_0)x + (\widehat{P}_2 - 2\widehat{P}_1 + \widehat{P}_0)x^2 + (\widehat{P}_3 - 2\widehat{P}_2 + \widehat{P}_1 - \widehat{P}_0)x^3 = \\ &= (j + 5\varepsilon + 4j\varepsilon) + (1 - \varepsilon - j\varepsilon)x + (\varepsilon + j\varepsilon)x^2 + (3j\varepsilon)x^3, \\ (1 - 2x + x^2 - x^3 + x^4)f_{\widehat{S}_n}(x) &= \widehat{S}_0 + (\widehat{S}_1 - 2\widehat{S}_0)x + (\widehat{S}_2 - 2\widehat{S}_1 + \widehat{S}_0)x^2 + (\widehat{S}_3 - 2\widehat{S}_2 + \widehat{S}_1 - \widehat{S}_0)x^3 = \\ &= (2j + 2\varepsilon + 5j\varepsilon + 4) + (\varepsilon - 2j - 6x - 4j\varepsilon)x + (3j - 2\varepsilon + 2)x^2 + (2\varepsilon - 4j + 8j\varepsilon + 7)x^3. \end{aligned}$$

Next, we give the exponential dual hyperbolic generating function of $\sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!}$ of the sequence \widehat{W}_n .

Lemma 2.4. Suppose that $f_{\widehat{W}_n}(x) = \sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!}$ is the exponential dual hyperbolic generating function of the generalized Pandita sequence $\{W_n\}$.

Then $\sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!}$ is given by

$$\sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!} = A_1 e^{\alpha x} \widehat{\alpha} + A_2 e^{\beta x} \widehat{\beta} + A_3 e^{\gamma x} \widehat{\gamma} + A_4 e^x \widehat{1}.$$

where $\widehat{\alpha}, \widehat{\beta}, \widehat{\gamma}, \widehat{1}$ are given as (2.5)-(2.8)

Proof. Using Binet's formula

$$W_n = A_1 \alpha^n + A_2 \beta^n + A_3 \gamma^n + A_4.$$

where A_1, A_2, A_3, A_4 are given in (1.4) we get

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} W_n \frac{x^n}{n!} + j \sum_{n=0}^{\infty} W_{n+1} \frac{x^n}{n!} + \varepsilon \sum_{n=0}^{\infty} W_{n+2} \frac{x^n}{n!} + j\varepsilon \sum_{n=0}^{\infty} W_{n+3} \frac{x^n}{n!} \\ &= \sum_{n=0}^{\infty} (A_1 \alpha^n + A_2 \beta^n + A_3 \gamma^n + A_4) \frac{x^n}{n!} + j \sum_{n=0}^{\infty} (A_1 \alpha^{n+1} + A_2 \beta^{n+1} + A_3 \gamma^{n+1} + A_4) \frac{x^n}{n!} \\ &\quad + \varepsilon \sum_{n=0}^{\infty} (A_1 \alpha^{n+2} + A_2 \beta^{n+2} + A_3 \gamma^{n+2} + A_4) \frac{x^n}{n!} + j\varepsilon \sum_{n=0}^{\infty} (A_1 \alpha^{n+3} + A_2 \beta^{n+3} + A_3 \gamma^{n+3} + A_4) \frac{x^n}{n!} \\ &= (A_1 e^{\alpha x} + A_2 e^{\beta x} + A_3 e^{\gamma x} + A_4 e^x) + j(A_1 \alpha e^{\alpha x} + A_2 \beta e^{\beta x} + A_3 \gamma e^{\gamma x} + A_4 e^x) \\ &\quad + \varepsilon(A_1 \alpha^2 e^{\alpha x} + A_2 \beta^2 e^{\beta x} + A_3 \gamma^2 e^{\gamma x} + A_4 e^x) + j\varepsilon(A_1 \alpha^3 e^{\alpha x} + A_2 \beta^3 e^{\beta x} + A_3 \gamma^3 e^{\gamma x} + A_4 e^x) \\ &= A_1 e^{\alpha x} (1 + j\alpha + \varepsilon\alpha^2 + j\varepsilon\alpha^3) + A_2 e^{\beta x} (1 + j\beta + \varepsilon\beta^2 + j\varepsilon\beta^3) \\ &\quad + A_3 e^{\gamma x} (1 + j\gamma + \varepsilon\gamma^2 + j\varepsilon\gamma^3) + A_4 e^x (1 + j + \varepsilon + j\varepsilon) \\ &= A_1 e^{\alpha x} \widehat{\alpha} + A_2 e^{\beta x} \widehat{\beta} + A_3 e^{\gamma x} \widehat{\gamma} + A_4 e^x \widehat{1} \end{aligned}$$

This proves (2.4). \square

The previous Lemma 2.4 gives the following results as particular examples.

Corollary 2.5. *Exponential dual hyperbolic generating function of Pandita and Pandita-Lucas numbers*

$$\begin{aligned} \text{a) } \sum_{n=0}^{\infty} \widehat{P}_n \frac{x^n}{n!} &= \frac{\alpha^3 e^{\alpha x} \widehat{\alpha}}{3\alpha - 2} + \frac{\beta^3 e^{\beta x} \widehat{\beta}}{3\beta - 2} + \frac{\gamma^3 e^{\gamma x} \widehat{\gamma}}{3\gamma - 2} - e^x \widehat{1}. \\ \text{b) } \sum_{n=0}^{\infty} \widehat{S}_n \frac{x^n}{n!} &= e^{\alpha x} \widehat{\alpha} + e^{\beta x} \widehat{\beta} + e^{\gamma x} \widehat{\gamma} + e^x \widehat{1}. \end{aligned}$$

3 OBTAINING BINET FORMULA FROM GENERATING FUNCTION

We next find Binet's formula generalized dual hyperbolic Pandita number $\{\widehat{W}_n\}$ by the use of generating function for \widehat{W}_n .

Theorem 3.1. *Binet's formula of generalized dual hyperbolic Pandita numbers*

$$\widehat{W}_n = \frac{q_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{q_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{q_3 \gamma^n}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{q_4 \delta^n}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}. \quad (3.1)$$

where

$$\begin{aligned} q_1 &= \widehat{W}_0 \alpha^3 + (\widehat{W}_1 - 2\widehat{W}_0) \alpha^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) \alpha - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3, \\ q_2 &= \widehat{W}_0 \beta^3 + (\widehat{W}_1 - 2\widehat{W}_0) \beta^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) \beta - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3, \\ q_3 &= \widehat{W}_0 \gamma^3 + (\widehat{W}_1 - 2\widehat{W}_0) \gamma^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) \gamma - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3, \\ q_4 &= \widehat{W}_0 \delta^3 + (\widehat{W}_1 - 2\widehat{W}_0) \delta^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) \delta - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3. \end{aligned}$$

Proof. Let

$$h(x) = x^4 - x^3 + x^2 - 2x + 1.$$

Then for some α, β, γ and δ we write

$$h(x) = (1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x).$$

i.e.,

$$x^4 - x^3 + x^2 - 2x + 1 = (1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x). \quad (3.2)$$

Hence $\frac{1}{\alpha}, \frac{1}{\beta}, \frac{1}{\gamma}$ and $\frac{1}{\delta}$ are the roots of $h(x)$. This gives α, β, γ and δ as the roots of

$$h\left(\frac{1}{x}\right) = \frac{1}{x^2} - \frac{2}{x} + \frac{1}{x^3} + \frac{1}{x^4} + 1 = 0.$$

This implies $x^4 - x^3 + x^2 - 2x + 1 = 0$. Now, by it follows that

$$\sum_{n=0}^{\infty} \widehat{W}_n x^n = \frac{(\widehat{W}_1 - \widehat{W}_0 - 2\widehat{W}_2 + \widehat{W}_3) x^3 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) x^2 + (\widehat{W}_1 - 2\widehat{W}_0) x + \widehat{W}_0}{(1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x)}.$$

Then we write

$$\begin{aligned} \frac{(\widehat{W}_1 - \widehat{W}_0 - 2\widehat{W}_2 + \widehat{W}_3) x^3 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) x^2 + (\widehat{W}_1 - 2\widehat{W}_0) x + \widehat{W}_0}{(1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x)} &= \frac{B_1}{(1 - \alpha x)} + \frac{B_2}{(1 - \beta x)} \\ &+ \frac{B_3}{(1 - \gamma x)} + \frac{B_4}{(1 - \delta x)}. \end{aligned} \quad (3.3)$$

So

$$\begin{aligned} &(\widehat{W}_1 - \widehat{W}_0 - 2\widehat{W}_2 + \widehat{W}_3) x^3 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) x^2 + (\widehat{W}_1 - 2\widehat{W}_0) x + \widehat{W}_0 \\ &= B_1(1 - \beta x)(1 - \gamma x)(1 - \delta x) + B_2(1 - \alpha x)(1 - \gamma x)(1 - \delta x) \\ &+ B_3(1 - \alpha x)(1 - \beta x)(1 - \delta x) + B_4(1 - \alpha x)(1 - \beta x)(1 - \gamma x). \end{aligned}$$

If we consider $x = \frac{1}{\alpha}$, we get $\widehat{W}_0 + \frac{1}{\alpha^2} (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) - \frac{1}{\alpha^3} (\widehat{W}_0 - \widehat{W}_1 + 2\widehat{W}_2 - \widehat{W}_3) + \frac{1}{\alpha} (\widehat{W}_1 - 2\widehat{W}_0) = -B_1 \left(\frac{1}{\alpha} \beta - 1 \right) \left(\frac{1}{\alpha} \gamma - 1 \right) \left(\frac{1}{\alpha} \delta - 1 \right)$.

This gives

$$\begin{aligned} B_1 &= \alpha^3 (\widehat{W}_0 + \frac{1}{\alpha^2} (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) + \frac{1}{\alpha^3} (\widehat{W}_1 - 5\widehat{W}_0 - 4\widehat{W}_2 + \widehat{W}_3) + \frac{1}{\alpha} (\widehat{W}_1 - 2\widehat{W}_0)) \\ &= \frac{\widehat{W}_0 \alpha^3 + (\widehat{W}_1 - 2\widehat{W}_0) \alpha^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2) \alpha - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)}. \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} B_2 &= \frac{\widehat{W}_0\beta^3 + (\widehat{W}_1 - 2\widehat{W}_0)\beta^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2)\beta - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)}, \\ B_3 &= \frac{\widehat{W}_0\gamma^3 + (\widehat{W}_1 - 2\widehat{W}_0)\gamma^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2)\gamma - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)}, \\ B_4 &= \frac{\widehat{W}_0\delta^3 + (\widehat{W}_1 - 2\widehat{W}_0)\delta^2 + (\widehat{W}_0 - 2\widehat{W}_1 + \widehat{W}_2)\delta - \widehat{W}_0 + \widehat{W}_1 - 2\widehat{W}_2 + \widehat{W}_3}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}. \end{aligned}$$

Thus (3.3) can be written as

$$\sum_{n=0}^{\infty} \widehat{W}_n x^n = B_1(1 - \alpha x)^{-1} + B_2(1 - \beta x)^{-1} + B_3(1 - \gamma x)^{-1} + B_4(1 - \delta x)^{-1}.$$

This gives

$$\sum_{n=0}^{\infty} \widehat{W}_n x^n = B_1 \sum_{n=0}^{\infty} \alpha^n x^n + B_2 \sum_{n=0}^{\infty} \beta^n x^n + B_3 \sum_{n=0}^{\infty} \gamma^n x^n + B_4 \sum_{n=0}^{\infty} \delta^n x^n = \sum_{n=0}^{\infty} (B_1 \alpha^n + B_2 \beta^n + B_3 \gamma^n + B_4 \delta^n) x^n.$$

Therefore, comparing coefficients on both sides of the above equality, we obtain

$$\widehat{W}_n = B_1 \alpha^n + B_2 \beta^n + B_3 \gamma^n + B_4 \delta^n.$$

The following identity establishes a relationship between the dual hyperbolic Pandita numbers and the Pandita-Lucas numbers.

Corollary 3.2. For all integers m, n the following identities holds:

$$\widehat{W}_{m+n} = P_{m-2} \widehat{W}_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5}) \widehat{W}_{n+2} + (P_{m-3} - P_{m-4}) \widehat{W}_{n+1} - \widehat{W}_n P_{m-3}.$$

Proof. First we assume that $m, n \geq 0$. The Theorem (3.2) can be proved by mathematical induction on m . If $m = 0$ we get

$$\widehat{W}_n = P_{-2} \widehat{W}_{n+3} + (P_{-4} - P_{-3} - P_{-5}) \widehat{W}_{n+2} + (P_{-3} - P_{-4}) \widehat{W}_{n+1} - \widehat{W}_n P_{-3}.$$

which is true since $P_{-2} = 0, P_{-1} = -1, P_{-4} = -1, P_{-5} = 0$. Assume that the equality holds for $m \leq k$. For $m = k + 1$, we get

$$\begin{aligned} \widehat{W}_{k+1+n} &= 2\widehat{W}_{n+k} - \widehat{W}_{n+k-1} + \widehat{W}_{n+k-2} - \widehat{W}_{n+k-3}, \\ &2(P_{m-2} \widehat{W}_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5}) \widehat{W}_{n+2} + (P_{m-3} - P_{m-4}) \widehat{W}_{n+1} - \widehat{W}_n P_{m-3}) \\ &- (P_{m-3} \widehat{W}_{n+3} + (P_{m-5} - P_{m-4} - P_{m-6}) \widehat{W}_{n+2} + (P_{m-4} - P_{m-5}) \widehat{W}_{n+1} - \widehat{W}_n P_{m-4}) \\ &+ (P_{m-4} \widehat{W}_{n+3} + (P_{m-6} - P_{m-5} - P_{m-7}) \widehat{W}_{n+2} + (P_{m-5} - P_{m-6}) \widehat{W}_{n+1} - \widehat{W}_n P_{m-5}) \\ &- (P_{m-5} \widehat{W}_{n+3} + (P_{m-7} - P_{m-6} - P_{m-8}) \widehat{W}_{n+2} + (P_{m-6} - P_{m-7}) \widehat{W}_{n+1} - \widehat{W}_n P_{m-6}). \end{aligned}$$

Consequently, by mathematical induction on m , this proves Theorem 3.2.

The other cases of m, n can be proved similarly for all integers m, n . \square

Taking $\widehat{W}_n = \widehat{P}_n$ or $\widehat{W}_n = \widehat{S}_n$ in above Theorem, respectively, we get:

Corollary 3.3.

$$\begin{aligned}\hat{P}_{m+n} &= P_{m-2}\hat{P}_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5})\hat{P}_{n+2} + (P_{m-3} - P_{m-4})\hat{P}_{n+1} - \hat{P}_n P_{m-3}, \\ \hat{S}_{m+n} &= P_{m-2}\hat{S}_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5})\hat{S}_{n+2} + (P_{m-3} - P_{m-4})\hat{S}_{n+1} - \hat{S}_n P_{m-3}.\end{aligned}$$

4 SIMSON'S FORMULAS

In this section, we present Simson's formula for the dual hyperbolic generalized Pandita numbers. This is a special case of [(Soykan, 2019c), Theorem 4.1].

Theorem 4.1. (Simpson's formula for dual hyperbolic generalized Pandita numbers) For all integers n we have,

$$\begin{vmatrix} \widehat{W}_{n+3} & \widehat{W}_{n+2} & \widehat{W}_{n+1} & \widehat{W}_n \\ \widehat{W}_{n+2} & \widehat{W}_{n+1} & \widehat{W}_n & \widehat{W}_{n-1} \\ \widehat{W}_{n+1} & \widehat{W}_n & \widehat{W}_{n-1} & \widehat{W}_{n-2} \\ \widehat{W}_n & \widehat{W}_{n-1} & \widehat{W}_{n-2} & \widehat{W}_{n-3} \end{vmatrix} = \begin{vmatrix} \widehat{W}_3 & \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 \\ \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} \\ \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} \\ \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} & \widehat{W}_{-3} \end{vmatrix} = (\widehat{W}_3 - 2\widehat{W}_2 + \widehat{W}_0)(\widehat{W}_3 - 2\widehat{W}_1 + \widehat{W}_0)(\widehat{W}_3^2 - \widehat{W}_2^2 + \widehat{W}_1^2 - \widehat{W}_0^2 - \widehat{W}_2\widehat{W}_3 - 2\widehat{W}_1\widehat{W}_3 + \widehat{W}_1\widehat{W}_2 + \widehat{W}_0\widehat{W}_3 + 2\widehat{W}_0\widehat{W}_2 - \widehat{W}_0\widehat{W}_1).$$

Proof. Using Theorem 2.1 it can be proved by using induction use [(Soykan, 2019c), Theorem 4.1]

From the Theorem 4.1 we get the following Corollary.

Corollary 4.2. For all integers n , the Simson's formulas of dual hyperbolic Pandita numbers and dual hyperbolic Pandita Lucas numbers are given as respectively

$$\begin{aligned}\text{a)} \quad & \begin{vmatrix} \hat{P}_{n+3} & \hat{P}_{n+2} & \hat{P}_{n+1} & \hat{P}_n \\ \hat{P}_{n+2} & \hat{P}_{n+1} & \hat{P}_n & \hat{P}_{n-1} \\ \hat{P}_{n+1} & \hat{P}_n & \hat{P}_{n-1} & \hat{P}_{n-2} \\ \hat{P}_n & \hat{P}_{n-1} & \hat{P}_{n-2} & \hat{P}_{n-3} \end{vmatrix} = 17 + 16j + 115\varepsilon + 113j\varepsilon. \\ \text{b)} \quad & \begin{vmatrix} \hat{S}_{n+3} & \hat{S}_{n+2} & \hat{S}_{n+1} & \hat{S}_n \\ \hat{S}_{n+2} & \hat{S}_{n+1} & \hat{S}_n & \hat{S}_{n-1} \\ \hat{S}_{n+1} & \hat{S}_n & \hat{S}_{n-1} & \hat{S}_{n-2} \\ \hat{S}_n & \hat{S}_{n-1} & \hat{S}_{n-2} & \hat{S}_{n-3} \end{vmatrix} = 452 + 655j + 1125\varepsilon - 126j\varepsilon.\end{aligned}$$

5 LINEAR SUMS

In this section, we give the summation formulas of the dual hyperbolic generalized Pandita numbers with positive and negative subscripts.

Now, we present the summation formulas of the generalized Pandita numbers.

Theorem 5.1. For the generalized Pandita numbers, we have the following formulas:

$$\begin{aligned}\text{(a)} \quad & \sum_{k=0}^n W_k = -(n+3)W_{n+3} + (n+4)W_{n+2} + (n+4)W_n + 3W_3 - 4W_2 - 3W_0. \\ \text{(b)} \quad & \sum_{k=0}^n W_{2k} = \frac{1}{3}(-3(n+2)W_{2n+2} + (3n+8)W_{2n+1} + 2W_{2n} + (3n+7)W_{2n-1} + 7W_3 - 8W_2 - W_1 - 6W_0). \\ \text{(c)} \quad & \sum_{k=0}^n W_{2k+1} = \frac{1}{3}(-(3n+4)W_{2n+2} + (3n+8)W_{2n+1} + W_{2n} + 3(n+2)W_{2n-1} + 6W_3 - 8W_2 + W_1 - 7W_0).\end{aligned}$$

Proof. For the proof, see [Soykan (2025), Theorem 3.12]. \square

Theorem 5.2. For the dual hyperbolic Pandita numbers, we have the following formulas:

- (a) $\sum_{k=0}^n \widehat{W}_k = -(n+3)\widehat{W}_{n+3} + (n+4)\widehat{W}_{n+2} + (n+4)\widehat{W}_n + 3\widehat{W}_3 - 4\widehat{W}_2 - 3\widehat{W}_0.$
- (b) $\sum_{k=0}^n \widehat{W}_{2k} = \frac{1}{3}(-3(n+2)\widehat{W}_{2n+2} + (3n+8)\widehat{W}_{2n+1} + 2\widehat{W}_{2n} + (3n+7)\widehat{W}_{2n-1} + 7\widehat{W}_3 - 8\widehat{W}_2 - \widehat{W}_1 - 6\widehat{W}_0).$
- (c) $\sum_{k=0}^n \widehat{W}_{2k+1} = \frac{1}{3}(-(3n+4)\widehat{W}_{2n+2} + (3n+8)\widehat{W}_{2n+1} + \widehat{W}_{2n} + 3(n+2)\widehat{W}_{2n-1} + 6\widehat{W}_3 - 8\widehat{W}_2 + \widehat{W}_1 - 7\widehat{W}_0).$

Proof. Use Theorem 5.1 and the definition of \widehat{W}_n . \square

As a special case of the theorem 5.2, we present the following Corollary.

Corollary 5.3. For $n \geq 0$, dual hyperbolic Pandita numbers have the following properties:

- (a) $\sum_{k=0}^n \widehat{P}_k = -(n+3)\widehat{P}_{n+3} + (n+4)\widehat{P}_{n+2} + (n+4)\widehat{P}_n + 1 - 5j\varepsilon - 2\varepsilon.$
- (b) $\sum_{k=0}^n \widehat{P}_{2k} = \frac{1}{3}(-3(n+2)\widehat{P}_{2n+2} + (3n+8)\widehat{P}_{2n+1} + 2\widehat{P}_{2n} + (3n+7)\widehat{P}_{2n-1} + 3j + \varepsilon - 3j\varepsilon + 4).$
- (c) $\sum_{k=0}^n \widehat{P}_{2k+1} = \frac{1}{3}(-(3n+4)\widehat{P}_{2n+2} + (3n+8)\widehat{P}_{2n+1} + \widehat{P}_{2n} + 3(n+2)\widehat{P}_{2n-1} + j - 3\varepsilon - 8j\varepsilon + 3).$

Corollary 5.4. For $n \geq 0$, dual hyperbolic Pandita Lucas numbers have the following properties.

- (a) $\sum_{k=0}^n \widehat{S}_k = -(n+3)\widehat{S}_{n+3} + (n+4)\widehat{S}_{n+2} + (n+4)\widehat{S}_n - 8j - 9\varepsilon - 10j\varepsilon - 5.$
- (b) $\sum_{k=0}^n \widehat{S}_{2k} = \frac{1}{3}(-3(n+2)\widehat{S}_{2n+2} + (3n+8)\widehat{S}_{2n+1} + 2\widehat{S}_{2n} + (3n+7)\widehat{S}_{2n-1} + -12j - 16\varepsilon - 15j\varepsilon - 7).$
- (c) $\sum_{k=0}^n \widehat{S}_{2k+1} = \frac{1}{3}(-(3n+4)\widehat{S}_{2n+2} + (3n+8)\widehat{S}_{2n+1} + \widehat{S}_{2n} + 3(n+2)\widehat{S}_{2n-1} + -16j - 15\varepsilon - 19j\varepsilon - 12).$

Next, we give the ordinary generating functions of some special cases of dual hyperbolic generalized Pandita numbers.

Theorem 5.5. The ordinary generating functions of the sequences \widehat{W}_{2n} , \widehat{W}_{2n+1} are given as follows:

- (a) $\sum_{n=0}^{\infty} \widehat{W}_{2n}x^n = \frac{\widehat{W}_2(x^3 + 3x^2 - x) + \widehat{W}_0(2x^2 + 2x - 1) - \widehat{W}_1(x^2 - x^3) - \widehat{W}_3(x^3 + 2x^2)}{-x^4 - x^3 + x^2 + 2x - 1}.$
- (b) $\sum_{n=0}^{\infty} \widehat{W}_{2n+1}x^n = \frac{\widehat{W}_0(x^3 + 2x^2) - \widehat{W}_3(x^3 + x^2 + x) - \widehat{W}_1(x^3 - 2x + 1) + \widehat{W}_2(2x^3 + x^2)}{-x^4 - x^3 + x^2 + 2x - 1}.$

Proof. Similary, the proof can be constructed as in [Theorem 2.2].

From the last Theorem, we have the following Corollary which gives sum formula of dual hyperbolic Pandita numbers (Take $\widehat{W}_n = \widehat{P}_n$ whit $\widehat{P}_0 = j + 2\varepsilon + 3j\varepsilon, \widehat{P}_1 = 2j + 3\varepsilon + 5j\varepsilon + 1, \widehat{P}_2 = 3j + 5\varepsilon + 8j\varepsilon + 2, \widehat{P}_3 = 5j + 8\varepsilon + 12j\varepsilon + 3$)

Corollary 5.6. For $n \geq 0$ dual hyperbolic Pandita numbers have the following properties.

- (a) $\sum_{n=0}^{\infty} \widehat{P}_{2n}x^n = \frac{(j + 5\varepsilon + 4j\varepsilon) + (1 - \varepsilon - j\varepsilon)x + (\varepsilon + j\varepsilon)x^2 + (3j\varepsilon)x^3}{1 - 2x + x^2 - x^3 + x^4},$
- (b) $\sum_{n=0}^{\infty} \widehat{P}_{2n+1}x^n = \frac{(2j + 2\varepsilon + 5j\varepsilon + 4) + (\varepsilon - 2j - 6x - 4j\varepsilon)x + (3j - 2\varepsilon + 2)x^2 + (2\varepsilon - 4j + 8j\varepsilon + 7)x^3}{1 - 2x + x^2 - x^3 + x^4}.$

6 MATRICES RELATED WITH DUAL HYPERBOLIC GENERALIZED PANDITA NUMBERS

In this section, using dual hyperbolic Pandita numbers, we give some matrices related to dual hyperbolic Pandita numbers.

We define the square matrix A of order 4 as

$$A = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

uch that $\det A = 1$. Note that

$$A^n = \begin{pmatrix} P_{n+1} & -P_n + P_{n-1} - P_{n-2} & P_n - P_{n-1} & -P_n \\ P_n & -P_{n-1} + P_{n-2} - P_{n-3} & P_{n-1} - P_{n-2} & -P_{n-1} \\ P_{n-1} & -P_{n-2} + P_{n-3} - P_{n-4} & P_{n-2} - P_{n-3} & -P_{n-2} \\ P_{n-2} & -P_{n-3} + P_{n-4} - P_{n-5} & P_{n-3} - P_{n-4} & -P_{n-3} \end{pmatrix}$$

for the proof see (Soykan, 2021e).

Then we give the following lemma.

Lemma 6.1. For $n \geq 0$ the following identity is true:

$$\begin{pmatrix} \widehat{W}_{n+3} \\ \widehat{W}_{n+2} \\ \widehat{W}_{n+1} \\ \widehat{W}_n \end{pmatrix} = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} \widehat{W}_3 \\ \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix}.$$

Proof. The identity(6.1) can be proved by mathematical induction on n . If $n = 0$ we obtain

$$\begin{pmatrix} \widehat{W}_3 \\ \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix} = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^0 \begin{pmatrix} \widehat{W}_3 \\ \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix}$$

which is true. We assume that the identity given holds for $n = k$. Thus the following identity is true

$$\begin{pmatrix} \widehat{W}_{k+3} \\ \widehat{W}_{k+2} \\ \widehat{W}_{k+1} \\ \widehat{W}_k \end{pmatrix} = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} \widehat{W}_3 \\ \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix}.$$

For $n = k + 1$, we get

$$\begin{aligned} \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^{k+1} \begin{pmatrix} \widehat{W}_3 \\ \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix} &= \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} \widehat{W}_3 \\ \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix} \\ &= \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \widehat{W}_{k+3} \\ \widehat{W}_{k+2} \\ \widehat{W}_{k+1} \\ \widehat{W}_k \end{pmatrix} \\ &= \begin{pmatrix} \widehat{W}_{k+4} \\ \widehat{W}_{k+3} \\ \widehat{W}_{k+2} \\ \widehat{W}_{k+1} \end{pmatrix}. \end{aligned}$$

Consequently, by mathematical induction on n , the proof completed. \square

We define

$$N_{\widehat{W}} = \begin{pmatrix} \widehat{W}_3 & \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 \\ \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} \\ \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} \\ \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} & \widehat{W}_{-3} \end{pmatrix}, \quad (6.1)$$

$$E_{\widehat{W}} = \begin{pmatrix} \widehat{W}_{n+3} & \widehat{W}_{n+2} & \widehat{W}_{n+1} & \widehat{W}_n \\ \widehat{W}_{n+2} & \widehat{W}_{n+1} & \widehat{W}_n & \widehat{W}_{n-1} \\ \widehat{W}_{n+1} & \widehat{W}_n & \widehat{W}_{n-1} & \widehat{W}_{n-2} \\ \widehat{W}_n & \widehat{W}_{n-1} & \widehat{W}_{n-2} & \widehat{W}_{n-3} \end{pmatrix}. \quad (6.2)$$

Now, we have the following theorem with $N_{\widehat{W}}$ and $E_{\widehat{W}}$

Theorem 6.2. Using $N_{\widehat{W}}$ and $E_{\widehat{W}}$, we get

$$A^n N_{\widehat{W}} = E_{\widehat{W}}.$$

Proof. Note that we get

$$\begin{aligned} A^n N_{\widehat{W}} &= \begin{pmatrix} P_{n+1} & -P_n + P_{n-1} - P_{n-2} & P_n - P_{n-1} & -P_n \\ P_n & -P_{n-1} + P_{n-2} - P_{n-3} & P_{n-1} - P_{n-2} & -P_{n-1} \\ P_{n-1} & -P_{n-2} + P_{n-3} - P_{n-4} & P_{n-2} - P_{n-3} & -P_{n-2} \\ P_{n-2} & -P_{n-3} + P_{n-4} - P_{n-5} & P_{n-3} - P_{n-4} & -P_{n-3} \end{pmatrix} \begin{pmatrix} \widehat{W}_3 & \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 \\ \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} \\ \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} \\ \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} & \widehat{W}_{-3} \end{pmatrix} \\ &= \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} \end{aligned}$$

where

$$\begin{aligned}
 a_{11} &= \widehat{W}_1(P_n - P_{n-1}) - \widehat{W}_2(P_n - P_{n-1} + P_{n-2}) - \widehat{W}_0P_n + W_3P_{n+1} = \widehat{W}_{n+3}, \\
 a_{12} &= \widehat{W}_0(P_n - P_{n-1}) - \widehat{W}_1(P_n - P_{n-1} + P_{n-2}) - P_n\widehat{W}_{-1} + \widehat{W}_2P_{n+1} = \widehat{W}_{n+2}, \\
 a_{13} &= \widehat{W}_{-1}(P_n - P_{n-1}) - \widehat{W}_0(P_n - P_{n-1} + P_{n-2}) - P_n\widehat{W}_{-2} + \widehat{W}_1P_{n+1} = \widehat{W}_{n+1}, \\
 a_{14} &= \widehat{W}_{-2}(P_n - P_{n-1}) - \widehat{W}_{-1}(P_n - P_{n-1} + P_{n-2}) - P_n\widehat{W}_{-3} + \widehat{W}_0P_{n+1} = \widehat{W}_n, \\
 a_{21} &= \widehat{W}_3P_n - \widehat{W}_2(P_{n-1} - P_{n-2} + P_{n-3}) + \widehat{W}(P_{n-1} - P_{n-2}) - \widehat{W}_0P_{n-1} = \widehat{W}_{n+2}, \\
 a_{22} &= \widehat{W}_2P_n - \widehat{W}_{-1}P_{n-1} - \widehat{W}_1(P_{n-1} - P_{n-2} + P_{n-3}) + \widehat{W}(P_{n-1} - P_{n-2}) = \widehat{W}_{n+1}, \\
 a_{23} &= \widehat{W}_{-1}(P_{n-1} - P_{n-2}) - \widehat{W}_{-2}P_{n-1} + \widehat{W}_1P_n - \widehat{W}_0(P_{n-1} - P_{n-2} + P_{n-3}) = \widehat{W}_n, \\
 a_{24} &= \widehat{W}_{-2}(P_{n-1} - P_{n-2}) - \widehat{W}_{-3}P_{n-1} + \widehat{W}_0P_n - \widehat{W}_{-1}(P_{n-1} - P_{n-2} + P_{n-3}) = \widehat{W}_{n-1}, \\
 a_{31} &= \widehat{W}_1(P_{n-2} - P_{n-3}) - \widehat{W}_2(P_{n-2} - P_{n-3} + P_{n-4}) - \widehat{W}_0P_{n-2} + \widehat{W}_3P_{n-1} = \widehat{W}_{n+1}, \\
 a_{32} &= \widehat{W}_0(P_{n-2} - P_{n-3}) - \widehat{W}_1(P_{n-2} - P_{n-3} + P_{n-4}) - \widehat{W}_{-1}P_{n-2} + \widehat{W}_2P_{n-1} = \widehat{W}_n, \\
 a_{33} &= \widehat{W}_{-1}(P_{n-2} - P_{n-3}) - \widehat{W}_{-2}P_{n-2} - \widehat{W}_0(P_{n-2} - P_{n-3} + P_{n-4}) + \widehat{W}_1P_{n-1} = \widehat{W}_{n-1}, \\
 a_{34} &= \widehat{W}_{-2}(P_{n-2} - P_{n-3}) - \widehat{W}_{-3}P_{n-2} - \widehat{W}_{-1}(P_{n-2} - P_{n-3} + P_{n-4}) + \widehat{W}_0P_{n-1} = \widehat{W}_{n-2}, \\
 a_{41} &= \widehat{W}_1(P_{n-3} - P_{n-4}) - \widehat{W}_2(P_{n-3} - P_{n-4} + P_{n-5}) - \widehat{W}_0P_{n-3} + \widehat{W}_3P_{n-2} = \widehat{W}_n, \\
 a_{42} &= \widehat{W}_0(P_{n-3} - P_{n-4}) - \widehat{W}_1(P_{n-3} - P_{n-4} + P_{n-5}) - \widehat{W}_{-1}P_{n-3} + \widehat{W}_2P_{n-2} = \widehat{W}_{n-1}, \\
 a_{43} &= \widehat{W}_{-1}(P_{n-3} - P_{n-4}) - \widehat{W}_{-2}P_{n-3} - \widehat{W}_0(P_{n-3} - P_{n-4} + P_{n-5}) + \widehat{W}_1P_{n-2} = \widehat{W}_{n-2}, \\
 a_{44} &= \widehat{W}_{-2}(P_{n-3} - P_{n-4}) - \widehat{W}_{-3}P_{n-3} - \widehat{W}_{-1}(P_{n-3} - P_{n-4} + P_{n-5}) + \widehat{W}_0P_{n-2} = \widehat{W}_{n-3}.
 \end{aligned}$$

Using the theorem (3.2) the proof is done. \square

By taking $\widehat{W}_n = \widehat{P}_n$ with $\widehat{P}_0, \widehat{P}_1, \widehat{P}_2, \widehat{P}_3$ in (6.1) and (6.2)
 $\widehat{W}_n = S_n$ with $\widehat{S}_0, \widehat{S}_1, \widehat{S}_2, \widehat{S}_3$ in (6.1) and (6.2)

respectively, we get:

$$\begin{aligned}
 N_{\widehat{P}} &= \begin{pmatrix} 5j+8\varepsilon+12j\varepsilon+3 & 3j+5\varepsilon+8j\varepsilon+2 & 2j+3\varepsilon+5j\varepsilon+1 & j+2\varepsilon+3j\varepsilon \\ 3j+5\varepsilon+8j\varepsilon+2 & 2j+3\varepsilon+5j\varepsilon+1 & j+2\varepsilon+3j\varepsilon & \varepsilon+2j\varepsilon \\ 2j+3\varepsilon+5j\varepsilon+1 & j+2\varepsilon+3j\varepsilon & \varepsilon+2j\varepsilon & -j\varepsilon \\ j+2\varepsilon+3j\varepsilon & \varepsilon+2j\varepsilon & -j\varepsilon & -1 \end{pmatrix}, \\
 E_{\widehat{P}} &= \begin{pmatrix} \widehat{P}_{n+3} & \widehat{P}_{n+2} & \widehat{P}_{n+1} & \widehat{P}_n \\ \widehat{P}_{n+2} & \widehat{P}_{n+1} & \widehat{P}_n & \widehat{P}_{n-1} \\ \widehat{P}_{n+1} & \widehat{P}_n & \widehat{P}_{n-1} & \widehat{P}_{n-2} \\ \widehat{P}_n & \widehat{P}_{n-1} & \widehat{P}_{n-2} & \widehat{P}_{n-3} \end{pmatrix}, \\
 N_{\widehat{S}} &= \begin{pmatrix} 6j+7\varepsilon+11j\varepsilon+5 & 5j+6\varepsilon+7j\varepsilon+2 & 2j+5\varepsilon+6j\varepsilon+2 & 2j+2\varepsilon+5j\varepsilon+4 \\ 5j+6\varepsilon+7j\varepsilon+2 & 2j+5\varepsilon+6j\varepsilon+2 & 2j+2\varepsilon+5j\varepsilon+4 & 4j+2\varepsilon+2j\varepsilon+1 \\ 2j+5\varepsilon+6j\varepsilon+2 & 2j+2\varepsilon+5j\varepsilon+4 & -4j+2\varepsilon+2j\varepsilon+1 & j+4\varepsilon+2j\varepsilon-1 \\ 2j+2\varepsilon+5j\varepsilon+4 & -4j+2\varepsilon+2j\varepsilon+1 & j+4\varepsilon+2j\varepsilon-1 & \varepsilon-j+4j\varepsilon+4 \end{pmatrix}, \\
 E_{\widehat{S}} &= \begin{pmatrix} \widehat{S}_{n+3} & \widehat{S}_{n+2} & \widehat{S}_{n+1} & \widehat{S}_n \\ \widehat{S}_{n+2} & \widehat{S}_{n+1} & \widehat{S}_n & \widehat{S}_{n-1} \\ \widehat{S}_{n+1} & S_n & \widehat{S}_{n-1} & \widehat{S}_{n-2} \\ \widehat{S}_n & \widehat{S}_{n-1} & \widehat{S}_{n-2} & \widehat{S}_{n-3} \end{pmatrix}.
 \end{aligned}$$

From Theorem (6.2), we can write the following corollary.

Corollary 6.3. *The following identities are hold:*

- a) $A^n N_{\hat{P}} = E_{\hat{P}}$.
- b) $A^n N_{\hat{S}} = E_{\hat{S}}$.

7 CONCLUSIONS

Recurrence relations define sequences where each term depends on previous ones. These sequences such as Fibonacci, Pell, Jacobsthal, Tribonacci, Padovan, Narayana's Cows, Leonardo, Tetranacci, and Pentanacci arise across fields including engineering, biology, mathematics, and physics. Below, we present their definitions with initial conditions using A_n notation and outline their real-world relevance.

- **Fibonacci Sequence:**

$$F_n = F_{n-1} + F_{n-2}, \quad F_0 = 0, \quad F_1 = 1$$

- **Pell Sequence:**

$$P_n = 2P_{n-1} + P_{n-2}, \quad P_0 = 0, \quad P_1 = 1$$

- **Jacobsthal Sequence:**

$$J_n = J_{n-1} + 2J_{n-2}, \quad J_0 = 0, \quad A_1 = 1$$

- **Tribonacci Sequence:**

$$T_n = T_{n-1} + T_{n-2} + T_{n-3}, \quad T_0 = 0, \quad T_1 = 1, \quad T_2 = 1$$

- **Padovan Sequence:**

$$P_n = P_{n-2} + P_{n-3}, \quad P_0 = P_1 = P_2 = 1$$

- **Narayana's Cows Sequence:**

$$N_n = N_{n-1} + N_{n-3}, \quad N_0 = N_1 = N_2 = 1$$

- **Leonardo Sequence:**

$$L_n = L_{n-1} + L_{n-2} + 1, \quad L_0 = 1, \quad L_1 = 1$$

- **Tetranacci Sequence:**

$$M_n = M_{n-1} + M_{n-2} + M_{n-3} + M_{n-4}, \quad M_0 = M_1 = M_2 = 0, \quad M_3 = 1$$

- **Pentanacci Sequence:**

$$P_n = P_{n-1} + P_{n-2} + P_{n-3} + P_{n-4} + P_{n-5}, \quad P_0 = P_1 = P_2 = P_3 = 0, \quad P_4 = 1$$

These sequences demonstrate how mathematical recursions extend into the fabric of our world whether designing structures, analyzing algorithms, modeling nature, or probing the quantum realm. Their recursive beauty continues to inspire both theoretical and practical exploration.

Next, we explore several real-world applications of recurrence relations across disciplines.

- Engineering
 - **Fibonacci:** Models recursive filters in control systems and signal processing.
 - **Padovan and Perrin:** Guide architectural proportions using the plastic number.
 - **Jacobsthal:** Applied in digital circuits for counting and encoding.
- Science
 - **Tribonacci and Tetranacci:** Simulate biological systems with delayed reproduction.
 - **Leonardo:** Reflect branching in plants and trees.
 - **Fibonacci and Narayana's Cows:** Describe phyllotaxis and seed arrangement in botany.
- Mathematics
 - **Recurrence Relations:** Analyze algorithms like mergesort and quicksort.
 - **Pell:** Solve Diophantine equations and approximate square roots with continued fractions.
 - **Jacobsthal and Padovan:** Used in tiling and combinatorics problems.
- Physics
 - **Fibonacci and Tribonacci:** Appear in wave interference and quantum systems.
 - **Pentanacci:** Used in recursive models of particle interactions and fractals.
 - **Padovan:** Linked to equilibrium modeling via the plastic constant.

In this study, we extend the classical framework to fourth-order recurrence systems by introducing the dual hyperbolic Pandita numbers, along with two distinguished subclasses. For these novel sequences, we derive Binet-type formulas, ordinary and exponential generating functions, and generalized Simson-type identities. Our analysis also encompasses closed-form summation formulas, algebraic properties, recurrence behaviors, and matrix-based representations.

Recognizing the theoretical depth and real-world utility of recurrence-based sequences, we first revisit the applications of second-order sequences to establish context. We then position our fourth-order generalizations as a natural progression within this broader mathematical landscape—offering new insights and powerful tools for modeling, analysis, and optimization in both pure and applied settings.

- For the applications of Gaussian Fibonacci and Gaussian Lucas numbers to Pauli Fibonacci and Pauli Lucas quaternions, see (Azak, 2022).
- For the application of Pell Numbers to the solutions of three-dimensional difference equation systems, see (Büyük and Taşkara, 2022).
- For the application of Jacobsthal numbers to special matrices, see (Vasanthi and Sivakumar, 2022).
- For the application of generalized k-order Fibonacci numbers to hybrid quaternions, see (Gül, 2022).
- For the applications of Fibonacci and Lucas numbers to Split Complex Bi-Periodic numbers, see (Yılmaz, 2022b).
- For the applications of generalized bivariate Fibonacci and Lucas polynomials to matrix polynomials, see (Yılmaz, 2022a).
- For the applications of generalized Fibonacci numbers to binomial sums, see (Ulutaş and Toy, 2022).
- For the application of generalized Jacobsthal numbers to hyperbolic numbers, see (Soykan and Taşdemir, 2022).
- For the application of generalized Fibonacci numbers to dual hyperbolic numbers, see (Soykan, 2021d).
- For the application of Laplace transform and various matrix operations to the characteristic

polynomial of the Fibonacci numbers, see (Deveci and Shannon, 2022).

- For the application of Generalized Fibonacci Matrices to Cryptography, see (Prasad and Mahato, 2022).
- For the application of higher order Jacobsthal numbers to quaternions, see (Özkan and Uysal, 2023).
- For the application of Fibonacci and Lucas Identities to Toeplitz-Hessenberg matrices, see (Goy and Shattuck, 2019).
- For the applications of Fibonacci numbers to lacunary statistical convergence, see (Bilgin, 2021).
- For the applications of Fibonacci numbers to lacunary statistical convergence in intuitionistic fuzzy normed linear spaces, see (Kişi and Tuzcuoglu, 2020).
- For the applications of Fibonacci numbers to ideal convergence on intuitionistic fuzzy normed linear spaces, see (Kişi and Debnath, 2022).
- For the applications of k -Fibonacci and k -Lucas numbers to spinors, see (Kumari et al., 2023).
- For the application of dual-generalized complex Fibonacci and Lucas numbers to Quaternions, see (Şentürk et al., 2022).
- For the application of special cases of Horadam numbers to Neutrosophic analysis see (Gökbaş et al., 2023).
- For the application of Hyperbolic Fibonacci numbers to Quaternions, see (Daşdemir, 2021).
- For the application of Pell numbers to Gaussian Hyperbolic numbers, see (Gökbaş, 2022).

In the following, we explore several applications of third-order recurrence sequences across various mathematical and applied contexts.

- For the applications of third order Jacobsthal numbers and Tribonacci numbers to quaternions, see (Cerdeira-Morales, 2017a) and (Cerdeira-Morales, 2017b), respectively.
- For the application of Tribonacci numbers to special matrices, see (Yilmaz and Taskara, 2014).
- For the applications of Padovan numbers and Tribonacci numbers to coding theory, see (Shtayat and Al-Kateeb, 2019) and (Basu and Das, 2014), respectively.

- For the application of Pell-Padovan numbers to groups, see (Deveci and Shannon, 2017).
- For the application of adjusted Jacobsthal-Padovan numbers to the exact solutions of some difference equations, see (Göçen, 2022).
- For the application of Gaussian Tribonacci numbers to various graphs, see (Sunitha and Sheriba, 2022).
- For the application of third-order Jacobsthal numbers to hyperbolic numbers, see (Dikmen and Altınsoy, 2022). For the application of Narayan numbers to finite groups see (Kuloğlu et al., 2022).
- For the application of generalized third-order Jacobsthal sequence to binomial transform, see (Soykan et al., 2022a).
- For the application of generalized Generalized Padovan numbers to Binomial Transform, see (Soykan et al., 2022b).
- For the application of generalized Tribonacci numbers to Gaussian numbers, see (Soykan et al., 2018).
- For the application of generalized Tribonacci numbers to Sedenions, see (Soykan et al., 2020a).
- For the application of Tribonacci and Tribonacci-Lucas numbers to matrices, see (Soykan, 2020).
- For the application of generalized Tribonacci numbers to circulant matrix, see (Soykan, 2021b).
- For the application of Tribonacci and Tribonacci-Lucas numbers to hybrinomials, see (Taşyurdu and Polat, 2021).
- For the application of hyperbolic Leonardo and hyperbolic Francois numbers to quaternions, see (Dişkaya et al., 2023).

In the following lists, we outline several applications of fourth-order recurrence sequences across theoretical and applied domains.

- For the application of Tetranacci and Tetranacci-Lucas numbers to quaternions, see (Soykan, 2019d).
- For the application of generalized Tetranacci numbers to Gaussian numbers, see (Soykan, 2019a).

- For the application of Tetranacci and Tetranacci-Lucas numbers to matrices, see (Soykan, 2019b).
- For the application of generalized Tetranacci numbers to binomial transform, see (Soykan, 2021c).

We now explore several applications of fifth-order sequences.

- For the application of Pentanacci numbers to matrices, see (Sivakumar and James, 2022).
- For the application of generalized Pentanacci numbers to quaternions, see (Soykan et al., 2020b).
- For the application of generalized Pentanacci numbers to binomial transform, see (Soykan, 2021a).

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Akar, M., Yüce, S., and Şahin, Ş. (2018). On the dual hyperbolic numbers and the complex hyperbolic numbers. *Journal of Computer Science & Computational Mathematics*, 8(1):1–6.
- Azak, A. Z. (2022). Pauli gaussian fibonacci and pauli gaussian lucas quaternions. *Mathematics*, 10:4655.
- Baez, J. (2002). The octonions. *Bulletin of the American Mathematical Society*, 39(2):145–205.
- Basu, M. and Das, M. (2014). Tribonacci matrices and a new coding theory. *Discrete Mathematics, Algorithms and Applications*, 6(1):1450008.
- Bilgin, N. G. (2021). Fibonacci lacunary statistical convergence of order γ in ifnls. *International Journal of Advances in Applied Mathematics and Mechanics*, 8(4):28–36.
- Biss, D. K., Dugger, D., and Isaksen, D. C. (2008). Large annihilators in cayley-dickson algebras. *Communications in Algebra*, 36(2):632–664.
- Bród, D., Liana, A., and Wloch, P. (2020). Two generalizations of dual-hyperbolic balancing numbers. *Symmetry*, 12(11):1866.
- Büyük, H. and Taşkara, N. (2022). On the solutions of three-dimensional difference equation systems via pell numbers. *European Journal of Science and Technology*, Special Issue 34:433–440.
- Cerda-Morales, G. (2017a). Identities for third order jacobsthal quaternions. *Advances in Applied Clifford Algebras*, 27(2):1043–1053.
- Cerda-Morales, G. (2017b). On a generalization of tribonacci quaternions. *Mediterranean Journal of Mathematics*, 14:239.
- Cheng, H. H. and Thompson, S. (1996). Dual polynomials and complex dual numbers for analysis of spatial mechanisms. In *Proc. of ASME 24th Biennial Mechanisms Conference*, pages 19–22, Irvine, CA.
- Cihan, A., Azak, A. Z., Güngör, M. A., and Tosun, M. (2019). A study on dual hyperbolic fibonacci and lucas numbers. *Analele Universitatii Ovidius Constanta - Seria Matematica*, 27(1):35–48.
- Cockle, J. (1849). On a new imaginary in algebra. *Philosophical Magazine, London-Dublin-Edinburgh*, 3(34):37–47.
- Daşdemir, A. (2021). On recursive hyperbolic fibonacci quaternions. *Communications in Advanced Mathematical Sciences*, 4(4):198–207.
- Deveci, O. and Shannon, A. G. (2017). Pell–padovan-circulant sequences and their applications. *Notes on Number Theory and Discrete Mathematics*, 23(3):100–114.
- Deveci, O. and Shannon, A. G. (2022). On recurrence results from matrix transforms. *Notes on Number Theory and Discrete Mathematics*, 28(4):589–592.

- Dikmen, C. M. (2025). A study on dual hyperbolic generalized leonardo numbers. *Asian Research Journal of Mathematics*, 21(6):143–165.
- Dikmen, C. M. and Altınsoy, M. (2022). On third order hyperbolic jacobsthal numbers. *Konuralp Journal of Mathematics*, 10(1):118–126.
- Dişkaya, O., Menken, H., and Catarino, P. M. M. C. (2023). On the hyperbolic leonardo and hyperbolic francois quaternions. *Journal of New Theory*, 42:74–85.
- Fjelstad, P. and Gal, S. (1998). n-dimensional hyperbolic complex numbers. *Advances in Applied Clifford Algebras*, 8(1):47–68.
- Göcen, M. (2022). The exact solutions of some difference equations associated with adjusted jacobsthal-padovan numbers. *Kırklareli University Journal of Engineering and Science*, 8(1):1–14.
- Gökbaş, H. (2022). Gaussian-hybrid numbers obtained from pell and pell-lucas sequences. *Journal of Advances in Mathematics and Computer Science*, 37(11):1–9.
- Gökbaş, H., Topal, S., and Smarandache, F. (2023). Neutrosophic number sequences: An introductory study. *International Journal of Neutrosophic Science (IJNS)*, 20(01):27–48.
- Goy, T. and Shattuck, M. (2019). Fibonacci and lucas identities from toeplitz-hessenberg matrices. *Applied and Applied Mathematics*, 14(2):699–715.
- Gül, K. (2022). Generalized k-order fibonacci hybrid quaternions. *Erzincan University Journal of Science and Technology*, 15(2):670–683.
- Hamilton, W. R. (1969). *Elements of Quaternions*. Chelsea Publishing Company, New York.
- Imaeda, K. and Imaeda, M. (2000). Sedenions: algebra and analysis. *Applied Mathematics and Computation*, 115:77–88.
- Kalça, F. and Soykan, Y. (2025). Gaussian numbers with generalized pandita numbers components. *Asian Journal of Advanced Research and Reports*, 19(7):32–56.
- Kantor, I. and Solodovnikov, A. (1989). *Hypercomplex Numbers*. Springer-Verlag, New York.
- Kiş, Ö. and Debnath, P. (2022). Fibonacci ideal convergence on intuitionistic fuzzy normed linear spaces. *Fuzzy Information and Engineering*, pages 1–13.
- Kiş, Ö. and Tuzcuoglu, I. (2020). Fibonacci lacunary statistical convergence in intuitionistic fuzzy normed linear spaces. *Journal of Progressive Research in Mathematics*, 16(3):3001–3007.
- Kuloğlu, B., Özkan, E., and Shannon, A. (2022). The narayana sequence in finite groups. *Fibonacci Quarterly*, 60(5):212–221.
- Kumari, M., Prasad, K., and Frontczak, R. (2023). On the k -fibonacci and k -lucas spinors. *Notes on Number Theory and Discrete Mathematics*, 29(2):322–335.
- Moreno, G. (1998). The zero divisors of the cayley-dickson algebras over the real numbers. *Boletín de la Sociedad Matemática Mexicana*, 3(4):13–28.
- Özkan, E. and Uysal, M. (2023). On quaternions with higher order jacobsthal numbers components. *Gazi University Journal of Science*, 36(1):336–347.
- Prasad, K. and Mahato, H. (2022). Cryptography using generalized fibonacci matrices with affine-hill cipher. *Journal of Discrete Mathematical Sciences & Cryptography*, 25(8-A):2341–2352.
- Şentürk, G. Y., Gürses, N., and Yüce, S. (2022). Construction of dual-generalized complex fibonacci and lucas quaternions. *Carpathian Mathematical Publications*, 14(2):406–418.
- Shtayat, J. and Al-Kateeb, A. (2019). An encoding-decoding algorithm based on padovan numbers. arXiv preprint.
- Sivakumar, B. and James, V. (2022). A notes on matrix sequence of pentanacci numbers and pentanacci cubes. *Communications in Mathematics and Applications*, 13(2):603–611.
- Sobczyk, G. (1995). The hyperbolic number plane. *The College Mathematics Journal*, 26(4):268–280.
- Soykan, Y. (2019a). Gaussian generalized tetranacci numbers. *Journal of Advances in Mathematics and Computer Science*, 31(3):1–21. Article no. JAMCS.48063.

- Soykan, Y. (2019b). Matrix sequences of tetranacci and tetranacci-lucas numbers. *International Journal of Advances in Applied Mathematics and Mechanics*, 7(2):57–69.
- Soykan, Y. (2019c). Simson identity of generalized m-step fibonacci numbers. *International Journal of Advanced Applied Mathematics and Mechanics*, 7(2):45–56.
- Soykan, Y. (2019d). Tetranacci and tetranacci-lucas quaternions. *Asian Research Journal of Mathematics*, 15(1):1–24. Article no. ARJOM.50749.
- Soykan, Y. (2019e). Tribonacci and tribonacci-lucas sedenions. *Mathematics*, 7(1):74.
- Soykan, Y. (2020). Matrix sequences of tribonacci and tribonacci-lucas numbers. *Communications in Mathematics and Applications*, 11(2):281–295.
- Soykan, Y. (2021a). Binomial transform of the generalized pentanacci sequence. *Asian Research Journal of Current Science*, 3(1):209–231.
- Soykan, Y. (2021b). Explicit euclidean norm, eigenvalues, spectral norm and determinant of circulant matrix with the generalized tribonacci numbers. *Earthline Journal of Mathematical Sciences*, 6(1):131–151.
- Soykan, Y. (2021c). On binomial transform of the generalized tetranacci sequence. *International Journal of Advances in Applied Mathematics and Mechanics*, 9(2):8–27.
- Soykan, Y. (2021d). On dual hyperbolic generalized fibonacci numbers. *Indian Journal of Pure and Applied Mathematics*.
- Soykan, Y. (2021e). Properties of generalized (r,s,t,u)-numbers. *Earthline Journal of Mathematical Sciences*, 5(2):297–327.
- Soykan, Y. (2023). Generalized pandita numbers. *International Journal of Mathematics, Statistics and Operations Research*, 3(1):107–123.
- Soykan, Y. (2025). Sums and generating functions of special cases of generalized tetranacci polynomials. *International Journal of Advances in Applied Mathematics and Mechanics*, 12(4):34–101.
- Soykan, Y., Gümüş, M., and Göcen, M. (2021). A study on dual hyperbolic generalized pell numbers. *Malaya Journal of Matematik*, 9(3):99–116.
- Soykan, Y., Okumuş, , and Taşdemir, E. (2020a). On generalized tribonacci sedenions. *Sarajevo Journal of Mathematics*, 16(1):103–122.
- Soykan, Y. and Taşdemir, E. (2022). A study on hyperbolic numbers with generalized jacobsthal numbers components. *International Journal of Nonlinear Analysis and Applications*, 13(2):1965–1981.
- Soykan, Y., Taşdemir, E., and Göcen, M. (2022a). Binomial transform of the generalized third-order jacobsthal sequence. *Asian-European Journal of Mathematics*, 15(12).
- Soykan, Y., Taşdemir, E., and Okumuş, (2022b). A study on binomial transform of the generalized padovan sequence. *Journal of Science and Arts*, 22(1):63–90.
- Soykan, Y., Taşdemir, E., and Okumuş, (2023). On dual hyperbolic numbers with generalized jacobsthal numbers components. *Indian Journal of Pure and Applied Mathematics*, 27(1):35–48.
- Soykan, Y., Taşdemir, E., Okumuş, , and Göcen, M. (2018). Gaussian generalized tribonacci numbers. *Journal of Progressive Research in Mathematics (JPRM)*, 14(2):2373–2387.
- Soykan, Y., Özmen, N., and Göcen, M. (2020b). On generalized pentanacci quaternions. *Tbilisi Mathematical Journal*, 13(4):169–181.
- Sunitha, K. and Sheriba, M. (2022). Gaussian tribonacci r-graceful labeling of some tree related graphs. *Ratio Mathematica*, 44:188–196.
- Taşyurdu, Y. and Polat, Y. E. (2021). Tribonacci and tribonacci-lucas hybrinomials. *Journal of Mathematics Research*, 13(5).
- Ulutaş, Y. T. and Toy, D. (2022). Some equalities and binomial sums about the generalized fibonacci number u_n . *Notes on Number Theory and Discrete Mathematics*, 28(2):252–260.
- Vasanthi, S. and Sivakumar, B. (2022). Jacobsthal matrices and their properties. *Indian Journal of Science and Technology*, 15(5):207–215.

- Yılmaz, B. and Soykan, Y. (2024). On dual hyperbolic guglielmo numbers. *Journal of Advances in Mathematics and Computer Science*, 39(4):37–61.
- Yılmaz, N. (2022a). The generalized bivariate fibonacci and lucas matrix polynomials. *Mathematica Montisnigri*, LIII:33–44.
- Yılmaz, N. (2022b). Split complex bi-periodic fibonacci and lucas numbers. *Communications Faculty of Science University of Ankara, Series A1 Mathematics and Statistics*, 71(1):153–164.
- Yılmaz, N. and Taskara, N. (2014). Tribonacci and tribonacci-lucas numbers via the determinants of special matrices. *Applied Mathematical Sciences*, 8(39):1947–1955.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://pr.sdiarticle5.com/review-history/141230>